

ELLIPTIC EQUATIONS WITH VERTICAL ASYMPTOTES IN THE NONLINEAR TERM

Par

LOUIS DUPAIGNE, AUGUSTO C. PONCE AND ALESSIO PORRETTA

Abstract. We study the existence of solutions of the nonlinear problem

$$(0.1) \quad \begin{cases} -\Delta u + g(u) = \mu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where μ is a bounded measure and g is a continuous nondecreasing function such that $g(0) = 0$. In this paper, we assume that the nonlinearity g satisfies

$$(0.2) \quad \lim_{t \uparrow 1} g(t) = +\infty.$$

Problem (0.1) need not have a solution for every measure μ . We prove that, given μ , there exists a “closest” measure μ^* for which (0.1) can be solved. We also explain how assumption (0.2) makes problem (0.1) different from the case where $g(t)$ is defined for every $t \in \mathbb{R}$.

1 Introduction

Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$, be a smooth bounded domain. In this paper, we are interested in the existence of solutions of the problem

$$(1.1) \quad \begin{cases} -\Delta u + g(u) = \mu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where μ is a bounded measure in Ω and $g : (-\infty, 1) \rightarrow \mathbb{R}$ is a continuous nondecreasing function such that $g(0) = 0$ and

$$(1.2) \quad \lim_{t \uparrow 1} g(t) = +\infty.$$

By a solution u of (1.1) we mean that $u \in L^1(\Omega)$, $u \leq 1$ a.e., $g(u) \in L^1(\Omega)$ and

$$-\int_{\Omega} u \Delta \zeta + \int_{\Omega} g(u) \zeta = \int_{\Omega} \zeta d\mu \quad \forall \zeta \in C^2(\overline{\Omega}), \zeta = 0 \text{ on } \partial\Omega.$$

In particular, $g(u) \in L^1(\Omega)$ implies that $u < 1$ a.e.

We observe that u , whenever it exists, is unique (see, e.g., [4]). It has been proved by Boccardo [2] (in the spirit of Brezis–Strauss [7]) that, for every $\mu \in L^1(\Omega)$, problem (1.1) has a solution. Moreover, Boccardo also shows that (1.1) has no solution if μ is a Dirac mass δ_a , with $a \in \Omega$. Consequently, we say that μ is a *good measure* (relative to g) if (1.1) has a solution u . We denote by $\mathcal{G}(g)$ the set of good measures associated to g .

Our goal in this paper is to investigate under what conditions on g and μ problem (1.1) admits a solution. We also point out to what extent assumption (1.2) makes this problem different from the case where g is a continuous function defined for every $t \in \mathbb{R}$, which was recently studied by Brezis–Marcus–Ponce [4].

We assume henceforth that, in addition to (1.2), g satisfies

$$(1.3) \quad g(t) = 0, \quad t \leq 0.$$

In particular, this implies that nonpositive measures are good for any g .

We denote by $\mathcal{M}(\Omega)$ the space of bounded Radon measures in Ω , equipped with its standard norm $\|\cdot\|_{\mathcal{M}}$. Given $\nu \in \mathcal{M}(\Omega)$, we say that ν is *diffuse* if $\nu(A) = 0$ for every Borel set $A \subset \Omega$ of zero H^1 -capacity (= Newtonian capacity). This capacity — which is denoted throughout this paper by “cap” — plays an important role in the study of problem (1.1).

The first consequence of (1.2) is that if (1.1) has a solution, then μ^+ is diffuse (see Corollary 2 in Section 2 below). The converse is not true; more precisely,

Theorem 1. *Given any g , there exists a diffuse measure $\mu \geq 0$ such that $\mu \notin \mathcal{G}(g)$.*

However, it turns out that every diffuse measure is good for *some* g (see Theorem 15).

For a fixed nonlinearity g , a natural question is to characterize the set of good measures associated to g . The next result gives a sufficient condition for a measure to be good.

Theorem 2. *Assume*

$$(1.4) \quad \limsup_{t \uparrow 1} \{(1-t)^{(2-\beta)/\beta} g(t)\} > 0$$

for some $0 < \beta < 2$. If $\mu^+ \ll \mathcal{H}^{N-2+\beta}$, then $\mu \in \mathcal{G}(g)$.

Here, \mathcal{H}^s denotes the s -dimensional Hausdorff measure of a set. By $\mu^+ \ll \mathcal{H}^s$, we mean that $\mu^+(A) = 0$ for every Borel set $A \subset \Omega$ such that $\mathcal{H}^s(A) = 0$. The

dimension $s = N - 2 + \beta$ in the statement of the theorem cannot be improved. In fact, given $\beta \in (0, 2)$, let

$$g(t) = \frac{1}{(1-t)^{(2-\beta)/\beta}} - 1, \quad t \in [0, 1).$$

For any $\alpha < N - 2 + \beta$, one can find a compact set $K_\alpha \subset \Omega$, with $\mathcal{H}^\alpha(K_\alpha) \in (0, \infty)$, such that if $\theta > 0$ is sufficiently large, then $\mu = \theta \mathcal{H}^\alpha \llcorner_{K_\alpha}$ is not good for g . This is easy to see if $\alpha \leq N - 2$, since in this case any compact set $K \subset \Omega$ such that $\mathcal{H}^\alpha(K) < \infty$ satisfies $\text{cap}(K) = 0$ (see, e.g., [8]); thus, by Corollary 2 in Section 2, μ is not good. In the remaining case, namely $N - 2 < \alpha < N - 2 + \beta$, the construction of K_α is rather delicate and is presented in Section 8 (see Theorem 18).

Even though the existence of solutions of problem (1.1) may fail for some diffuse measures (by Theorem 1), $L^1(\Omega)$ is not the largest set where (1.1) has a solution for any g . For instance, let $\mu \in \mathcal{M}(\Omega)$ be such that $v \leq 1$ a.e., where v is the unique solution of

$$(1.5) \quad \begin{cases} -\Delta v = \mu & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Then μ is good for every g (see Proposition 7 in Section 7). The converse is also true if μ^+ is singular with respect to the Lebesgue measure in \mathbb{R}^N . In fact, we have the following.

Theorem 3. *Let $\mu \in \mathcal{M}(\Omega)$ be such that μ^+ is singular. Then $\mu \in \mathcal{G}(g)$ for every g if and only if $v \leq 1$ a.e., where v is given by (1.5).*

The characterization of the set of all measures in $\mathcal{M}(\Omega)$ which are good for every g is given in Section 7.

Our method in the study of problem (1.1) starts with a standard procedure, which consists in approximating g with bounded continuous functions defined on the whole \mathbb{R} . More precisely, let (g_n) be a sequence of bounded functions $g_n : \mathbb{R} \rightarrow \mathbb{R}$ which are continuous, nondecreasing and satisfy the following conditions:

$$(1.6) \quad 0 \leq g_1(t) \leq g_2(t) \leq \dots, \quad t \in \mathbb{R},$$

$$(1.7) \quad g_n(t) \rightarrow g(t), \quad t < 1,$$

and

$$(1.8) \quad g_n(t) \rightarrow +\infty, \quad t \geq 1.$$

Since each g_n is bounded, there exists a unique solution u_n of

$$(1.9) \quad \begin{cases} -\Delta u_n + g_n(u_n) = \mu & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega. \end{cases}$$

Passing to the limit as n tends to infinity we get the following result.

Proposition 1. *Given any $\mu \in \mathcal{M}(\Omega)$, then $u_n \downarrow u^*$ in Ω as $n \uparrow +\infty$, where u^* is the largest subsolution of (1.1). Moreover, we have*

$$(1.10) \quad \|g(u^*)\|_{L^1} \leq \|\mu\|_{\mathcal{M}}$$

and

$$(1.11) \quad \left| \int_{\Omega} u^* \Delta \zeta \right| \leq 2 \|\mu\|_{\mathcal{M}} \|\zeta\|_{L^\infty} \quad \forall \zeta \in C_0^2(\bar{\Omega}).$$

Here,

$$C_0^2(\bar{\Omega}) = \left\{ \zeta \in C^2(\bar{\Omega}) : \zeta = 0 \text{ on } \partial\Omega \right\}.$$

In the spirit of [4], we define the *reduced measure* μ^* as

$$\mu^* = -\Delta u^* + g(u^*)$$

and study the properties of μ^* . First of all, since u^* is the largest subsolution of (1.1), μ^* is well-defined, independently of the sequence (g_n) . Note that $\mu^* \leq \mu$; moreover, μ is a good measure if and only if $\mu = \mu^*$.

Theorem 4. *For every $\mu \in \mathcal{M}(\Omega)$, there exist Borel sets $\Sigma_1, \Sigma_2 \subset \Omega$ such that*

$$(1.12) \quad \Sigma_1 \subset [u^* = 1], \quad \text{cap}(\Sigma_2) = 0, \quad \text{and} \quad (\mu - \mu^*)(\Omega \setminus (\Sigma_1 \cup \Sigma_2)) = 0.$$

Note that, in the previous statement, the set $[u^* = 1]$ is well-defined up to sets of zero H^1 -capacity. Indeed, any function $v \in L^1(\Omega)$ such that $\Delta v \in \mathcal{M}(\Omega)$ admits a unique cap-quasicontinuous representative \tilde{v} (see, e.g., [1]); henceforth, we always identify v and \tilde{v} . We recall that \tilde{v} is cap-quasicontinuous if, for every $\varepsilon > 0$, there exists an open set $\omega_\varepsilon \subset \Omega$ such that $\text{cap}(\omega_\varepsilon) < \varepsilon$ and $\tilde{v}|_{\Omega \setminus \omega_\varepsilon}$ is continuous.

We remark that, in Theorem 4, both sets Σ_1 and Σ_2 have zero Lebesgue measure, so we have the following.

Corollary 1. *For any measure μ , we have*

$$(\mu^*)_{\mathbf{a}} = \mu_{\mathbf{a}},$$

where “a” denotes the absolutely continuous part with respect to the Lebesgue measure.

In view of Theorem 4, if μ is diffuse and $\mu([u^* = 1]) = 0$, then it follows that $\mu^* = \mu$; hence μ is good. We use this idea to prove Theorem 2; in this case, the main effort is thus to estimate the $(N - 2 + \beta)$ -Hausdorff measure of the set $[u^* = 1]$.

This kind of estimate, which has an interest of its own, is given by Theorem 12 in Section 4. In Section 5, we present another approach, based on energy estimates; in this case, the “smallness” of $[u^* = 1]$ is given in terms of (Sobolev) capacities.

The next result says that μ^* is the “best approximation” of μ in the class of good measures relative to g .

Theorem 5. *For $\mu \in \mathcal{M}(\Omega)$,*

$$(1.13) \quad \|\mu - \mu^*\|_{\mathcal{M}} = \min_{\nu \in \mathcal{G}} \|\mu - \nu\|_{\mathcal{M}}.$$

In addition, μ^ is the unique good measure for which the minimum in (1.13) is attained.*

We recall that when the function g is defined for every $t \in \mathbb{R}$, it has been shown in [4] that μ^* is the largest good measure $\leq \mu$. In that case, the characterization of μ^* given in Theorem 5 is then a straightforward consequence. An important difference in our case is that there exist measures μ for which the set $\{\lambda \in \mathcal{G}(g) : \lambda \leq \mu\}$ has *no* largest element (see Proposition 9 in Section 9). Thus, the fact that μ^* is the unique measure which achieves the minimum in (1.13) needs a direct proof, which is more delicate.

Finally, two further differences with the case studied in [4] are worth mentioning. When $g(t)$ is defined for every $t \in \mathbb{R}$, the set \mathcal{G} of good measures is convex, and the mapping $\mu \mapsto \mu^*$ is a contraction. As is shown in Section 9 below, these properties are no longer true when g satisfies (1.2). In fact, for any such g ,

- (a) \mathcal{G} is *not* convex;
- (b) the mapping $\mu \mapsto \mu^*$ is *not* a contraction.

We emphasize that throughout this paper we assume that Ω is a domain of \mathbb{R}^N with $N \geq 2$. The case of dimension $N = 1$ is different and has been studied by Vázquez [20]. We recall that in this case, every measure is diffuse — since $\text{cap}(\{x\}) > 0$ for every x — and the solutions of (1.1) are Lipschitz continuous. In [20], Vázquez proves that

- (a') if $\int_0^1 g = +\infty$, then every $\mu \in \mathcal{M}(\Omega)$ is good;
- (b') if $\int_0^1 g < +\infty$ and $\mu \in \mathcal{M}(\Omega)$ satisfies $\|\mu^+\|_{\mathcal{M}} \leq 2\sqrt{2} \left(\int_0^1 g \right)^{1/2}$, then μ is good.

These two results have no counterpart when $N \geq 2$. According to Theorem 1 above, for any g , there exists a diffuse measure $\mu \geq 0$ such that μ is not good. As we see in Section 8, such μ can be chosen so that $\varepsilon\mu$ is not good for *any* $\varepsilon > 0$.

The plan of this paper is the following:

1. Introduction
2. Proofs of Proposition 1 and Theorem 4
3. The reduced measure is the closest good measure
4. Proof of Theorem 2
5. Capacitary estimates related to problem (1.1)
6. Every diffuse measure is good for some g
7. Measures which are good for every g
8. How to construct diffuse measures which are not good
9. Further properties of μ^* and \mathcal{G}

2 Proofs of Proposition 1 and Theorem 4

We start by recalling (see, e.g., [16]) that every measure μ can be uniquely decomposed as

$$\mu = \mu_d + \mu_c,$$

where μ_d is diffuse and μ_c is concentrated on a set of zero capacity. In particular, μ is diffuse if and only if $\mu_c = 0$.

A useful characterization of measures which are diffuse is given by

Theorem 6 ([3, 17]). *Let $\mu \in \mathcal{M}(\Omega)$. Then μ is diffuse if and only if*

$$\mu \in L^1(\Omega) + H^{-1}(\Omega).$$

The next two results are used frequently in the sequel.

Theorem 7 ([6]). *Let $v \in L^1(\Omega)$ be such that $\Delta v \in \mathcal{M}(\Omega)$. Then*

$$\Delta v^+ \in \mathcal{M}_{\text{loc}}(\Omega)$$

and

$$(2.1) \quad (\Delta v^+)_d \geq \chi_{[v \geq 0]} (\Delta v)_d \quad \text{in } \Omega,$$

$$(2.2) \quad (-\Delta v^+)_c = (-\Delta v)_c^+ \quad \text{in } \Omega.$$

Moreover, if $v \geq 0$ a.e., then

$$(2.3) \quad (\Delta v)_d \geq 0 \quad \text{in } [v = 0].$$

Theorem 8 ([14]). *Let $v \in L^1(\Omega)$ be such that $\Delta v \in \mathcal{M}(\Omega)$. If $v \geq 0$ a.e., then*

$$(2.4) \quad (\Delta v)_c \leq 0 \quad \text{in } \Omega.$$

As a result, we get a necessary condition for (1.1) to admit a solution.

Corollary 2. *If μ is good, then μ^+ is diffuse.*

Proof. Applying Theorem 8 to $v = 1 - u$, we get

$$\mu_c = (-\Delta u)_c = (\Delta v)_c \leq 0.$$

Thus $\mu^+ = (\mu_d)^+ = (\mu^+)_d$, and so μ^+ is diffuse. \square

Let us also recall that given $\nu \in \mathcal{M}(\Omega)$, there exists a unique function $v \in L^1(\Omega)$ which satisfies

$$-\int_{\Omega} v \Delta \zeta = \int_{\Omega} \zeta d\nu \quad \forall \zeta \in C_0^2(\overline{\Omega}).$$

This function is called Stampacchia's solution of the problem (see [19])

$$(2.5) \quad \begin{cases} -\Delta v = \nu & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega, \end{cases}$$

and coincides with the notion of "renormalized solution" introduced in [9]. In particular, Theorems 2.33 and 10.1 of [9] provide the following useful result.

Theorem 9 ([9]). *Let v be the unique solution of (2.5). Let $\Phi \in W^{2,\infty}(\mathbb{R})$ be such that $\text{supp } \Phi''$ is compact. Then*

$$\Delta \Phi(v) = \Phi'(v)(\Delta v)_d + \Phi''(v)|\nabla v|^2 - \Phi'(+\infty)(\Delta v)_c^- + \Phi'(-\infty)(\Delta v)_c^+ \quad \text{in } \Omega.$$

Here we denote by $\Phi'(\pm\infty)$ the limit of Φ' as $|x| \rightarrow \pm\infty$.

The proof of Proposition 1 follows the same lines as in [4]. We present the proof for the convenience of the reader.

Proof of Proposition 1. Let u_n be the solution of (1.9). Since $g_n \leq g_{n+1}$, by a comparison principle (see, e.g., [4, Appendix B]) we have $u_n \geq u_{n+1}$. Hence we define u^* such that

$$u_n \downarrow u^* \quad \text{a.e. in } \Omega.$$

Standard estimates imply that

$$(2.6) \quad \|g_n(u_n)\|_{L^1} \leq \|\mu\|_{\mathcal{M}};$$

thus,

$$\left| \int_{\Omega} u_n \Delta \zeta \right| = \left| \int_{\Omega} \zeta d\mu - \int_{\Omega} g_n(u_n) \zeta \right| \leq 2\|\mu\|_{\mathcal{M}} \|\zeta\|_{L^\infty} \quad \forall \zeta \in C_0^2(\overline{\Omega}).$$

Clearly, $u^* \in L^1(\Omega)$ and (u_n) converges strongly to u^* in $L^1(\Omega)$. Moreover, it follows from (2.6) that $u^* \leq 1$ a.e. Then, using Fatou's lemma, we deduce from the previous estimates that

- (i) $g(u^*) \in L^1(\Omega)$ and (1.10) holds;
- (ii) $\Delta u^* \in \mathcal{M}(\Omega)$ and $\|\Delta u^*\|_{\mathcal{M}} \leq 2\|\mu\|_{\mathcal{M}}$.

Finally, let v be any subsolution of (1.1), i.e., $v \in L^1(\Omega)$, $v \leq 1$ a.e., $g(v) \in L^1(\Omega)$ and

$$-\int_{\Omega} v \Delta \zeta + \int_{\Omega} g(v) \zeta \leq \int_{\Omega} \zeta d\mu \quad \forall \zeta \in C_0^2(\overline{\Omega}), \zeta \geq 0 \text{ in } \Omega.$$

Since $g_n \leq g$, we have

$$-\Delta v + g_n(v) \leq -\Delta v + g(v) \leq \mu = -\Delta u_n + g_n(u_n) \quad \text{in } [C_0^2(\overline{\Omega})]^*,$$

which yields $v \leq u_n$ a.e. Passing to the limit, we deduce that $v \leq u^*$. This proves that u^* is the largest subsolution of (1.1). \square

Let

$$(2.7) \quad \mu^* = -\Delta u^* + g(u^*) \quad \text{in } \mathcal{D}'(\Omega).$$

In view of Proposition 1, $\mu^* \in \mathcal{M}(\Omega)$. The reduced measure μ^* is uniquely determined by the weak* limit of $g_n(u_n)$ in $\mathcal{M}(\Omega)$. Indeed, comparing (2.7) with (1.9), and using that $u_n \rightarrow u^*$ in $L^1(\Omega)$, we obtain the following.

Lemma 1. *Let (u_n) be the sequence defined in Proposition 1. Then*

$$(2.8) \quad g_n(u_n) \xrightarrow{*} g(u^*) + (\mu - \mu^*) \quad \text{weak* in } \mathcal{M}(\Omega).$$

Note that since $\Delta u^* \in \mathcal{M}(\Omega)$, the function u^* admits a unique cap-quasicontinuous representative, which we use henceforth as our standard choice. In particular, we remark that the set $[u^* = 1]$ is uniquely defined up to sets of zero capacity.

The main ingredient in the proof of Theorem 4 is

Proposition 2. *Let u^* be given by Proposition 1 and let μ^* be the reduced measure defined in (2.7). Then*

$$(2.9) \quad 0 \leq \mu - \mu^* \leq (\mu_d)_{[u^*=1]} + \mu_c^+ \quad \text{in } \Omega.$$

In particular,

$$(2.10) \quad \begin{cases} (\mu^*)_d = \mu_d & \text{in } [u^* < 1], \\ (\mu^*)_d \geq 0 & \text{in } [u^* = 1], \\ (\mu^*)_c = -(\mu_c)^- & \text{in } \Omega, \\ (\mu^*)^- = \mu^- & \text{in } \Omega. \end{cases}$$

Proof.

Step 1. Proof of (2.9).

Given $\delta > 0$, define the function $\theta_\delta(s) = \min \{1, \frac{1}{\delta}(s - 1 + 2\delta)^+\}$. Applying Theorem 9 with $v = u_n$ and $\Phi_\delta(s) = \int_0^s \theta_\delta(\xi) d\xi$, we get for any $\zeta \in C_0^2(\overline{\Omega})$, $\zeta \geq 0$ in Ω ,

$$(2.11) \quad \int_{\Omega} g_n(u_n) \theta_\delta(u_n) \zeta dx \leq \int_{\Omega} \theta_\delta(u_n) \zeta d\mu_d + \int_{\Omega} \zeta d\mu_c^+ + \int_{\Omega} \Phi_\delta(u_n) \Delta \zeta dx,$$

which yields

$$\int_{[u_n > 1-\delta]} g_n(u_n) \zeta dx \leq \int_{\Omega} \theta_\delta(u_n) \zeta d\mu_d + \int_{\Omega} \zeta d\mu_c^+ + \int_{\Omega} \Phi_\delta(u_n) \Delta \zeta dx.$$

Since (u_n^+) is bounded in $L^\infty(\Omega)$ and (Δu_n) is bounded in $\mathcal{M}(\Omega)$, the sequence $(\theta_\delta(u_n))$ is bounded in $H_0^1(\Omega)$, converges weakly to $\theta_\delta(u^*)$ in $H_0^1(\Omega)$ and weak* in $L^\infty(\Omega)$. Moreover, since $\mu_d \in L^1(\Omega) + H^{-1}(\Omega)$ (by Theorem 6), we have

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \theta_\delta(u_n) \zeta d\mu_d = \int_{\Omega} \theta_\delta(u^*) \zeta d\mu_d.$$

Thus,

$$\limsup_{n \rightarrow +\infty} \int_{[u_n > 1-\delta]} g_n(u_n) \zeta dx \leq \int_{\Omega} \theta_\delta(u^*) \zeta d\mu_d + \int_{\Omega} \zeta d\mu_c^+ + \int_{\Omega} \Phi_\delta(u^*) \Delta \zeta dx.$$

Clearly, by dominated convergence we have, for a.e. $\delta > 0$,

$$\lim_{n \rightarrow +\infty} \int_{[u_n \leq 1-\delta]} g_n(u_n) \zeta \, dx = \int_{[u^* \leq 1-\delta]} g(u^*) \zeta \, dx.$$

Therefore,

$$\begin{aligned} & \limsup_{n \rightarrow +\infty} \int_{\Omega} g_n(u_n) \zeta \, dx \\ & \leq \int_{[u^* \leq 1-\delta]} g(u^*) \zeta \, dx + \int_{\Omega} \theta_{\delta}(u^*) \zeta \, d\mu_d + \int_{\Omega} \zeta \, d\mu_c^+ + \int_{\Omega} \Phi_{\delta}(u^*) \Delta \zeta \, dx, \end{aligned}$$

for a.e. $\delta > 0$. Since $u^* < 1$ a.e., $\Phi_{\delta}(u^*) \rightarrow 0$ a.e. By dominated convergence, as $\delta \rightarrow 0$ we obtain

$$\limsup_{n \rightarrow +\infty} \int_{\Omega} g_n(u_n) \zeta \, dx \leq \int_{\Omega} g(u^*) \zeta \, dx + \int_{[u^*=1]} \zeta \, d\mu_d + \int_{\Omega} \zeta \, d\mu_c^+.$$

Comparing with (2.8), we get

$$\mu - \mu^* \leq (\mu_d)_{[u^*=1]} + \mu_c^+.$$

Clearly, by Fatou's lemma, $\mu^* \leq \mu$. We thus obtain (2.9).

Step 2. Proof of (2.10).

From (2.9), we deduce immediately that

$$(2.12) \quad \begin{aligned} (\mu^*)_d &= \mu_d \quad \text{in } [u^* < 1], \\ (\mu^*)_d &\geq 0 \quad \text{in } [u^* = 1]. \end{aligned}$$

Since $\mu_d \geq (\mu^*)_d$, (2.12) yields

$$(2.13) \quad (\mu_d)^- = (\mu^*)_d^- \quad \text{in } \Omega.$$

On the other hand, by (2.9),

$$\mu_c - (\mu^*)_c \leq \mu_c^+ \quad \text{in } \Omega;$$

that is,

$$(\mu^*)_c \geq -\mu_c^- \quad \text{in } \Omega.$$

Note that $\mu^* \leq \mu$ and $(\mu^*)_c \leq 0$ (by Corollary 2); thus

$$(\mu^*)_c \leq -\mu_c^- \quad \text{in } \Omega.$$

We conclude that

$$(2.14) \quad (\mu^*)_c = -\mu_c^- \quad \text{in } \Omega.$$

Assertion (2.10) then follows from (2.12)–(2.14). \square

As a consequence of the previous result, we have the

Proof of Theorem 4. Let $\Sigma_1 = [u^* = 1]$ and let $\Sigma_2 \subset \Omega$ be such that $\text{cap}(\Sigma_2) = 0$ and $\mu_c^+(\Omega \setminus \Sigma_2) = 0$. With this choice, (1.12) follows immediately from (2.9). \square

As a corollary of (2.10), we also have

Corollary 3. *Let $\mu \in \mathcal{M}(\Omega)$. If $\mu \geq 0$, then $\mu^* \geq 0$.*

We give an alternative characterization of u^* in the next result.

Proposition 3. *For every $\mu \in \mathcal{M}(\Omega)$, u^* is the unique solution of the following problem:*

$$(2.15) \quad \begin{cases} v \in W_0^{1,1}(\Omega), & v \leq 1 \text{ a.e.}, & \Delta v \in \mathcal{M}(\Omega), & g(v) \in L^1(\Omega), \\ (-\Delta v)_d + g(v) = \mu_d & \text{in } [v < 1], \\ (-\Delta v)_d \leq \mu_d & \text{in } [v = 1], \\ (-\Delta v)_c = -\mu_c^- & \text{in } \Omega. \end{cases}$$

Proof. From (2.7) and (2.10), it follows that u^* is indeed a solution of (2.15). We now prove that the solution of (2.15) is unique. Assume that v_1, v_2 both satisfy (2.15) and consider the function $w = (v_1 - v_2)^+$. First observe that Δw is a measure and, by (2.3),

$$(2.16) \quad (\Delta w)_d \geq 0 \quad \text{in } [v_1 \leq v_2].$$

By (2.15),

$$(2.17) \quad (\Delta v_1)_d \geq g(v_1) - \mu_d \quad \text{in } \Omega.$$

On the other hand, since $[v_1 > v_2] \subset [v_2 < 1]$, we have

$$(2.18) \quad (\Delta v_2)_d = g(v_2) - \mu_d \quad \text{in } [v_1 > v_2].$$

Thus, by (2.17)–(2.18),

$$(2.19) \quad (\Delta w)_d \geq [\Delta(v_1 - v_2)]_d \geq g(v_1) - g(v_2) \geq 0 \quad \text{in } [v_1 > v_2].$$

It follows from (2.16) and (2.19) that

$$(2.20) \quad (\Delta w)_d \geq 0 \quad \text{in } \Omega.$$

Since, by (2.2),

$$(-\Delta w)_c = [-\Delta(v_1 - v_2)]_c^+ \quad \text{in } \Omega,$$

we get

$$(2.21) \quad (-\Delta w)_c = [(-\Delta v_1)_c + (\Delta v_2)_c]^+ = 0 \quad \text{in } \Omega.$$

From (2.20)–(2.21), we obtain that

$$\Delta w \geq 0 \quad \text{in } \Omega.$$

Since w vanishes on $\partial\Omega$, we have $v_1 \leq v_2$ a.e. in Ω (see, e.g., [4, Proposition B.1]). Reversing the roles of the two functions, we finally obtain that $v_1 = v_2$. \square

Until now, we have studied problem (1.1) by approximating the nonlinearity g using a sequence (g_n) , with μ fixed. Another possible approach is to fix g and to approximate μ by $\rho_n * \mu$, where (ρ_n) is a sequence of mollifiers. More precisely, $\rho_n * \mu$ is given by

$$(\rho_n * \mu)(x) = \int_{\Omega} \rho_n(x-y) d\mu(y) \quad \forall x \in \Omega.$$

It turns out that the sequences of solutions in both cases converge to the same limit.

Theorem 10. *Let $\mu \in \mathcal{M}(\Omega)$. For each $n \geq 1$, let v_n be the solution of*

$$(2.22) \quad \begin{cases} -\Delta v_n + g(v_n) = \rho_n * \mu & \text{in } \Omega, \\ v_n = 0 & \text{on } \partial\Omega. \end{cases}$$

Then, $v_n \rightarrow u^$ in $L^1(\Omega)$, where u^* is the function given by Proposition 1.*

Proof. By standard estimates, we have

$$\|g(v_n)\|_{L^1(\Omega)} \leq \|\rho_n * \mu\|_{\mathcal{M}(\Omega)} \leq \|\mu\|_{\mathcal{M}(\Omega)}.$$

Thus, Δv_n is bounded in $L^1(\Omega)$ and there exist $v \in L^1(\Omega)$ and $\nu \in \mathcal{M}(\Omega)$ such that, for a subsequence (still denoted (v_n)), we have

$$\begin{aligned} v_n &\rightarrow v \quad \text{strongly in } L^1(\Omega) \text{ and a.e.} \\ g(v_n) &\xrightarrow{*} g(v) + \nu \quad \text{weak* in } \mathcal{M}(\Omega). \end{aligned}$$

By Fatou's lemma, $\nu \geq 0$. Moreover, it follows that v satisfies

$$(2.23) \quad -\Delta v + g(v) = \mu - \nu \quad \text{in } \Omega.$$

We now follow the outline of the proof of Proposition 2. Take $\theta_\delta(v_n)\zeta$ as a test function in (2.22), where $\zeta \in C_0^2(\overline{\Omega})$. We get the analogue of (2.11), namely

$$\int_{\Omega} g(v_n) \theta_\delta(v_n) \zeta dx \leq \int_{\Omega} \theta_\delta(v_n) (\rho_n * \mu_d) \zeta dx + \int_{\Omega} (\rho_n * \mu_c^+) \zeta dx + \int_{\Omega} \Phi_\delta(v_n) \Delta \zeta dx.$$

As in Step 1 of Proposition 2, we obtain, as n tends to infinity, that

$$(2.24) \quad \nu \leq (\mu_d)_{[v=1]} + \mu_c^+ \quad \text{in } \Omega.$$

Using (2.23)–(2.24), we obtain that v is a solution of (2.15) (see Step 2 of Proposition 2). From the uniqueness result of Proposition 3, we conclude that $v = u^*$. In particular, the whole sequence (v_n) converges to u^* . \square

3 The reduced measure is the closest good measure

We start with the following simple result.

Proposition 4. *Let $\mu \in \mathcal{M}(\Omega)$. If $\mu \in \mathcal{G}(g)$ and $\nu \leq \mu$, then $\nu \in \mathcal{G}(g)$.*

Proof. Let (u_n) be the sequence of functions satisfying (1.9). By standard estimates (see, e.g., [4]), we have

$$\int_{\Omega} |g_n(u_n) - g_n(u)| \leq \int_{\Omega} |g(u) - g_n(u)|.$$

Thus,

$$\int_{\Omega} |g_n(u_n) - g(u)| \leq 2 \int_{\Omega} |g(u) - g_n(u)| \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

We conclude that $g_n(u_n) \rightarrow g(u)$ in $L^1(\Omega)$. Let (v_n) be the sequence associated to ν . By comparison, $\nu \leq \mu$ implies $v_n \leq u_n$ a.e.; thus $g_n(v_n) \leq g_n(u_n)$ a.e. Applying the dominated convergence theorem, we conclude that $g_n(v_n) \rightarrow g(v^*)$ in $L^1(\Omega)$. We then deduce that v^* is the solution of (1.1) with data ν , and so ν is a good measure. \square

We also have the following.

Lemma 2. *For every $\mu, \nu \in \mathcal{M}(\Omega)$,*

$$(3.1) \quad \int_{\Omega} |g(u^*) - g(v^*)| + \|(\mu - \mu^*) - (\nu - \nu^*)\|_{\mathcal{M}} \leq \|\mu - \nu\|_{\mathcal{M}},$$

where u^*, v^* are the solutions of (1.1) with respect to μ^*, ν^* , resp.

Proof. Let v_n denote the solution of

$$\begin{cases} -\Delta v_n + g_n(v_n) = \nu & \text{in } \Omega, \\ v_n = 0 & \text{on } \partial\Omega. \end{cases}$$

By Lemma 1, we get

$$g_n(u_n) - g_n(v_n) \xrightarrow{*} g(u^*) - g(v^*) + (\mu - \mu^*) - (\nu - \nu^*) \quad \text{weak}^* \text{ in } \mathcal{M}(\Omega).$$

Since $\mu - \mu^*$ and $\nu - \nu^*$ are both singular with respect to Lebesgue measure (see Corollary 1), we have

$$\|g(u^*) - g(v^*) + (\mu - \mu^*) - (\nu - \nu^*)\|_{\mathcal{M}} = \int_{\Omega} |g(u^*) - g(v^*)| + \|(\mu - \mu^*) - (\nu - \nu^*)\|_{\mathcal{M}}.$$

On the other hand,

$$\|g_n(u_n) - g_n(v_n)\|_{L^1} \leq \|\mu - \nu\|_{\mathcal{M}} \quad \forall n \geq 1.$$

Thus,

$$\begin{aligned} \int_{\Omega} |g(u^*) - g(v^*)| + \|(\mu - \mu^*) - (\nu - \nu^*)\|_{\mathcal{M}} &\leq \liminf_{n \rightarrow +\infty} \|g_n(u_n) - g_n(v_n)\|_{L^1} \\ &\leq \|\mu - \nu\|_{\mathcal{M}}, \end{aligned}$$

which gives (3.1). □

A simple consequence is the following result.

Theorem 11. *For every $\mu, \nu \in \mathcal{M}(\Omega)$,*

$$(3.2) \quad \|\mu^* - \nu^*\|_{\mathcal{M}} \leq 2\|\mu - \nu\|_{\mathcal{M}}.$$

Proof. Let $\mu, \nu \in \mathcal{M}(\Omega)$. By Lemma 2,

$$\|(\mu - \mu^*) - (\nu - \nu^*)\|_{\mathcal{M}} \leq \|\mu - \nu\|_{\mathcal{M}}.$$

Applying the triangle inequality, we obtain (3.2). □

We now present the

Proof of Theorem 5. We split the proof into two steps.

Step 1. Proof of (1.13).

Given $\nu \in \mathcal{G}$, we have $\nu = \nu^*$. It then follows from Lemma 2 that

$$(3.3) \quad \int_{\Omega} |g(u^*) - g(v)| + \|\mu - \mu^*\|_{\mathcal{M}} \leq \|\mu - \nu\|_{\mathcal{M}},$$

where v is the solution of (1.1) with measure ν . In particular,

$$\|\mu - \mu^*\|_{\mathcal{M}} \leq \|\mu - \nu\|_{\mathcal{M}},$$

which gives (1.13).

Step 2. μ^* is the unique good measure which achieves the minimum in (1.13).

We now assume that $\nu \in \mathcal{G}$ satisfies

$$(3.4) \quad \|\mu - \nu\|_{\mathcal{M}} = \|\mu - \mu^*\|_{\mathcal{M}}.$$

By (3.3), we have

$$\int_{\Omega} |g(u^*) - g(v)| = 0.$$

Thus,

$$(3.5) \quad g(u^*) = g(v) \quad \text{a.e.}$$

We next observe that $\nu \leq \mu$. In fact, note that

$$(3.6) \quad \inf \{\mu, \nu\} = \mu - (\mu - \nu)^+.$$

Moreover, by Proposition 4, $\inf \{\mu, \nu\} \leq \nu$ implies that $\inf \{\mu, \nu\}$ is also a good measure. It then follows from (3.6) and the minimality of ν that

$$\|\mu - \nu\|_{\mathcal{M}} \leq \|\mu - \inf \{\mu, \nu\}\|_{\mathcal{M}} = \|(\mu - \nu)^+\|_{\mathcal{M}}.$$

Therefore, $(\mu - \nu)^- = 0$; in other words, $\nu \leq \mu$. In particular, ν is a subsolution of (1.1), so that $\nu \leq u^*$ a.e. by Proposition 1.

We now split the proof into two cases.

Case 1. $\text{cap}([u^* = 1]) = 0$.

By Theorem 4, this implies $(\mu - \mu^*)_{\text{d}} = 0$. Thus,

$$\nu_{\text{d}} \leq \mu_{\text{d}} = (\mu^*)_{\text{d}}.$$

On the other hand, since $\nu \leq u^*$ a.e., it follows from Theorem 8 that

$$\nu_{\text{c}} = (-\Delta \nu)_{\text{c}} \leq (-\Delta u^*)_{\text{c}} = (\mu^*)_{\text{c}}.$$

We conclude that

$$\nu \leq \mu^* \leq \mu.$$

By (3.4), we must have $\nu = \mu^*$.

Case 2. $\text{cap}([u^* = 1]) > 0$.

We first show that $u^* = v$ on a set of positive Lebesgue measure. By way of contradiction, suppose that $v < u^*$ a.e. Let $\alpha_0, \beta_0 \in [0, 1]$ be such that $\alpha_0 < \beta_0$ and g is increasing on $[\alpha_0, \beta_0]$. Since (3.5) holds and $v < u^*$ a.e., the set $[\alpha_0 < u^* < \beta_0]$ has zero Lebesgue measure. Let

$$w = \min \{\beta_0, \max \{\alpha_0, u^*\}\} - \alpha_0.$$

Thus, $w \in H_0^1(\Omega)$ and w assumes only the values 0 and $\beta_0 - \alpha_0$. We conclude that $w = 0$ a.e. In other words, $u^* \leq \alpha_0$ a.e. Since $\text{cap}([u^* = 1]) > 0$, we get a contradiction.

We now proceed with the proof of Case 2. Given $\varepsilon > 0$, let $\alpha, \beta \in (1 - \varepsilon, 1)$, $\alpha < \beta$, be such that g is increasing in $[\alpha, \beta]$. Let $\Phi_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that $\Phi_\varepsilon(t) = t$ if $t \leq \alpha$, $\Phi_\varepsilon(t) = 1$ if $t \geq \beta$ and $\Phi'_\varepsilon(t) \geq 0$, $\forall t \in \mathbb{R}$. We now establish the following.

Claim. *For every $\varepsilon > 0$, we have*

$$(3.7) \quad -\Delta[\Phi_\varepsilon(u^*) - \Phi_\varepsilon(v)] \geq 0 \quad \text{in } \mathcal{D}'(\Omega).$$

Proof. In fact, by Theorem 9,

$$(3.8) \quad \begin{aligned} [\Delta\Phi_\varepsilon(u^*)]_d &= \Phi'_\varepsilon(u^*)(\Delta u^*)_d + \Phi''_\varepsilon(u^*)|\nabla u^*|^2 \\ &= \Phi'_\varepsilon(u^*)[g(u^*) - (\mu^*)_d] + \Phi''_\varepsilon(u^*)|\nabla u^*|^2 \end{aligned}$$

and, similarly,

$$(3.9) \quad [\Delta\Phi_\varepsilon(v)]_d = \Phi'_\varepsilon(v)[g(v) - \nu_d] + \Phi''_\varepsilon(v)|\nabla v|^2.$$

By construction of Φ_ε , we have

$$(3.10) \quad \Phi'_\varepsilon(u^*) = \Phi'_\varepsilon(v) \quad \text{a.e.}$$

This is clear if $v \leq u^* \leq \alpha$ or $\beta \leq v \leq u^*$. Finally, if $\alpha < u^*$ and $v < \beta$, then $u^* = v$ a.e., since g is increasing in $[\alpha, \beta]$ and $g(u^*) = g(v)$ a.e. We conclude that (3.10) holds.

By (3.5) and (3.10), we then have

$$(3.11) \quad \Phi'_\varepsilon(u^*)g(u^*) - \Phi'_\varepsilon(v)g(v) = 0 \quad \text{a.e.}$$

Note that

$$\Phi''_\varepsilon(u^*) = \Phi''_\varepsilon(v) \quad \text{a.e.}$$

In addition, on the set where $\Phi''_\varepsilon(u^*) \neq 0$, we have $u^* = v$ a.e., so that

$$\nabla u^* = \nabla v \quad \text{a.e. in } [\Phi''_\varepsilon(u^*) \neq 0].$$

Thus,

$$(3.12) \quad \Phi''_\varepsilon(u^*)|\nabla u^*|^2 - \Phi''_\varepsilon(v)|\nabla v|^2 = 0 \quad \text{a.e.}$$

Finally, since $\Phi'_\varepsilon(1) = 0$ and $(\mu^*)_d = \mu_d$ on the set $[u^* < 1]$ (by Theorem 4), we have

$$\Phi'_\varepsilon(u^*)(\mu^*)_d = \Phi'_\varepsilon(u^*)\mu_d \quad \text{in } \Omega.$$

Moreover, $\Phi'_\varepsilon \geq 0$ and $\nu \leq \mu$ imply

$$\Phi'_\varepsilon(v)\nu_d \leq \Phi'_\varepsilon(v)\mu_d \quad \text{in } \Omega.$$

Therefore,

$$(3.13) \quad \Phi'_\varepsilon(u^*)(\mu^*)_d - \Phi'_\varepsilon(v)\nu_d \geq [\Phi'_\varepsilon(u^*) - \Phi'_\varepsilon(v)]\mu_d = 0 \quad \text{in } \Omega.$$

Subtracting (3.9) from (3.8) and then applying (3.11)–(3.13), we conclude that

$$(3.14) \quad -\left(\Delta[\Phi_\varepsilon(u^*) - \Phi_\varepsilon(v)]\right)_d \geq 0 \quad \text{in } \Omega.$$

On the other hand, since $u^* \geq v$ a.e., we have $\Phi_\varepsilon(u^*) - \Phi_\varepsilon(v) \geq 0$ a.e. It then follows from Theorem 8 that

$$(3.15) \quad -\left(\Delta[\Phi_\varepsilon(u^*) - \Phi_\varepsilon(v)]\right)_c \geq 0 \quad \text{in } \Omega.$$

Combining (3.14) and (3.15), we obtain (3.7). This concludes the proof of the claim. \square

According to the previous claim, the function $\Phi_\varepsilon(u^*) - \Phi_\varepsilon(v)$ is superharmonic. Moreover, since it is nonnegative and $\Phi_\varepsilon(u^*) = \Phi_\varepsilon(v)$ a.e. on a set of positive (Lebesgue) measure, we deduce from the strong maximum principle (see [1]; see also [5]) that

$$\Phi_\varepsilon(u^*) = \Phi_\varepsilon(v) \quad \text{a.e. in } \Omega.$$

Since this holds true for every $\varepsilon > 0$, we let $\varepsilon \downarrow 0$ and conclude that $u^* = v$ a.e. Thus, $\mu^* = \nu$. The proof of Theorem 5 is complete. \square

4 Proof of Theorem 2

In order to establish Theorem 2, we assume the next result, which will be proved afterwards.

Theorem 12. *Let $v \in L^1(\Omega)$, $v \leq 1$ a.e., be such that $\Delta v \in \mathcal{M}(\Omega)$. Assume g satisfies (1.4) for some $0 < \beta < 2$. If $g(v) \in L^1(\Omega)$, then*

$$(4.1) \quad \mathcal{H}^{N-2+\beta}([v = 1]) = 0.$$

Proof of Theorem 2. Clearly, it suffices to establish the theorem for $\mu \geq 0$. Let u^* be the function given by Proposition 1. Since $\Delta u^* \in \mathcal{M}(\Omega)$ and $g(u^*) \in L^1(\Omega)$, it follows from Theorem 12 that

$$\mathcal{H}^{N-2+\beta}([u^* = 1]) = 0.$$

By assumption, we have $\mu \ll \mathcal{H}^{N-2+\beta}$. Thus, μ is diffuse and

$$\mu([u^* = 1]) = 0.$$

It follows from Theorem 4 that $\mu^* = \mu$. In other words, $\mu \in \mathcal{G}$. This concludes the proof of Theorem 2. \square

We split the proof of Theorem 12 into two cases: $0 < \beta < 1$ and $1 \leq \beta < 2$. We first consider the case $0 < \beta < 1$. An important ingredient is the following.

Lemma 3. *Let $\nu \in \mathcal{M}(\Omega)$ and let v be the solution of*

$$(4.2) \quad \begin{cases} -\Delta v = \nu & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Given $0 < \beta < 1$ and $k \geq 1$, there exists a Borel set $A_k \subset \Omega$ such that

$$(4.3) \quad |v(x) - v(y)| \leq Ck |x - y|^\beta \quad \forall x, y \in \Omega \setminus A_k$$

and

$$(4.4) \quad \mathcal{H}_\infty^{N-2+\beta}(A_k) \leq \frac{C}{k} \|\nu\|_{\mathcal{M}},$$

for some constant $C > 0$ independent of k .

Given $\alpha \geq 0$, the Hausdorff content $\mathcal{H}_\infty^\alpha$ of a Borel set $A \subset \mathbb{R}^N$ is defined as

$$\mathcal{H}_\infty^\alpha(A) = \inf \left\{ \sum_i r_i^\alpha : A \subset \bigcup_i B_{r_i}(x_i) \right\},$$

where the infimum is taken over all coverings of A with balls $B_{r_i}(x_i)$ of radii r_i . Note that we make no restriction on the size of such balls. In particular, for every bounded set A , $\mathcal{H}_\infty^\alpha(A) < \infty$. It is easy to see that

$$(4.5) \quad \mathcal{H}_\infty^\alpha(A) = 0 \quad \text{if and only if} \quad \mathcal{H}^\alpha(A) = 0.$$

Proof of Lemma 3. By linearity, it suffices to establish the lemma for $\nu \geq 0$. Let

$$(4.6) \quad A_k = \left\{ x \in \Omega : \nu(B_r(x)) \geq k r^{N-2+\beta} \text{ for some } r > 0 \right\}.$$

(Here, ν is viewed as a measure in \mathbb{R}^N such that $\nu(\mathbb{R}^N \setminus \Omega) = 0$).

We claim that (4.3) and (4.4) hold for A_k . We begin by establishing (4.4). For each $x \in A_k$, let $r_x > 0$ be such that

$$\nu(B_{r_x}(x)) \geq k r_x^{N-2+\beta}.$$

Clearly, $(B_{5r_x}(x))_{x \in A_k}$ is a covering of A_k . Applying Vitali's covering lemma, we may extract a subcovering $(B_{5r_i}(x_i))$ of A_k such that the balls $B_{r_i}(x_i)$ are all disjoint. We then have

$$\begin{aligned} \mathcal{H}_\infty^{N-2+\beta}(A_k) &\leq \sum_i (5r_i)^{N-2+\beta} \\ &= C \sum_i r_i^{N-2+\beta} \\ &\leq \frac{C}{k} \sum_i \nu(B_{r_i}(x_i)) = \frac{C}{k} \nu\left(\bigcup_i B_{r_i}(x_i)\right) \leq \frac{C}{k} \|\nu\|_{\mathcal{M}}. \end{aligned}$$

This is precisely (4.4). We now turn to the proof of (4.3). We follow closely the argument presented in [8]. For simplicity, we assume $N \geq 3$; the case $N = 2$ follows the same lines.

Clearly, it suffices to prove (4.3) for the function

$$w(x) = \frac{1}{N(N-2)\omega_N} \int_\Omega \frac{d\nu(z)}{|z-x|^{N-2}} \quad \forall x \in \Omega,$$

where $\omega_N = |B_1|$ is the measure of the unit ball in \mathbb{R}^N . It is not difficult to see that w can be rewritten as (see, e.g., [18, Lemma 2])

$$w(x) = \frac{1}{N\omega_N} \int_0^\infty \frac{\nu(B_s(x))}{s^{N-1}} ds.$$

Given $x, y \in \Omega \setminus A_k$, let $\delta = |x - y|$. We then write

$$\begin{aligned} (4.7) \quad w(x) - w(y) &= \frac{1}{N\omega_N} \int_0^\infty \left[\nu(B_s(x)) - \nu(B_s(y)) \right] \frac{ds}{s^{N-1}} \\ &= \frac{1}{N\omega_N} \left\{ \int_0^{2\delta} + \int_{2\delta}^\infty \right\}. \end{aligned}$$

Since $x, y \notin A_k$,

$$\nu(B_s(x)), \nu(B_s(y)) \leq k s^{N-2+\beta} \quad \forall s > 0.$$

We then have

$$(4.8) \quad \int_0^{2\delta} \leq \int_0^{2\delta} \nu(B_s(x)) \frac{ds}{s^{N-1}} \leq k \int_0^{2\delta} \frac{ds}{s^{1-\beta}} = Ck \delta^\beta.$$

On the other hand, for $s \geq 2\delta$, $B_{s-\delta}(x) \subset B_s(y)$; thus,

$$\begin{aligned} \int_{2\delta}^{\infty} &\leq \int_{2\delta}^{\infty} \left[\nu(B_s(x)) - \nu(B_{s-\delta}(x)) \right] \frac{ds}{s^{N-1}} \\ &\leq \int_{\delta}^{\infty} \nu(B_s(x)) \left\{ \frac{1}{s^{N-1}} - \frac{1}{(s+\delta)^{N-1}} \right\} ds. \end{aligned}$$

Since

$$\frac{1}{s^{N-1}} - \frac{1}{(s+\delta)^{N-1}} \leq C \frac{\delta}{s^N} \quad \forall s \geq \delta,$$

we obtain

$$(4.9) \quad \int_{2\delta}^{\infty} \leq C\delta \int_{\delta}^{\infty} \nu(B_s(x)) \frac{ds}{s^N} \leq Ck\delta \int_{\delta}^{\infty} \frac{ds}{s^{2-\beta}} \leq Ck\delta^{\beta}.$$

It follows from (4.7)–(4.9) that

$$w(x) - w(y) \leq Ck\delta^{\beta} = Ck|x-y|^{\beta}.$$

Switching the roles between x and y , we conclude that w satisfies (4.3). Since $v-w$ is a harmonic function, v also verifies (4.3). The proof of the lemma is complete. \square

Given a Borel set $A \subset \mathbb{R}^N$, let

$$\Theta^*(x, A) = \limsup_{t \rightarrow 0} \frac{|A \cap B_t(x)|}{|B_t(x)|},$$

where $|\cdot|$ denotes Lebesgue measure in \mathbb{R}^N . This function gives the density of points of A which are close to x . Clearly, $0 \leq \Theta^*(x, A) \leq 1$.

Lemma 4. *Given a Borel set $A \subset \mathbb{R}^N$, let*

$$(4.10) \quad F = \{x \in \mathbb{R}^N : \Theta^*(x, A) \geq 1/4\}.$$

Then, for every $0 \leq \alpha \leq N$,

$$(4.11) \quad \mathcal{H}_{\infty}^{\alpha}(F) \leq C \mathcal{H}_{\infty}^{\alpha}(A)$$

for some $C > 0$ depending on N and α .

Proof. If $\alpha = 0$, the conclusion is clear. Assume $\alpha > 0$. Given $\varepsilon > 0$, let $(B_{r_i}(x_i))$ be a covering of A such that

$$(4.12) \quad \sum_i r_i^{\alpha} \leq \mathcal{H}_{\infty}^{\alpha}(A) + \varepsilon.$$

Let

$$F_1 = F \cap \left[\bigcup_i B_{2r_i}(x_i) \right] \quad \text{and} \quad F_2 = F \setminus \left[\bigcup_i B_{2r_i}(x_i) \right].$$

Clearly,

$$(4.13) \quad \mathcal{H}_\infty^\alpha(F_1) \leq \sum_i (2r_i)^\alpha \leq 2^\alpha \left[\mathcal{H}_\infty^\alpha(A) + \varepsilon \right].$$

We now prove a similar estimate for $\mathcal{H}_\infty^\alpha(F_2)$. Since (4.10) holds, for each $y \in F_2$, one can find $s_y > 0$ sufficiently small so that

$$(4.14) \quad |A \cap B_{s_y/2}(y)| \geq \frac{1}{8} |B_{s_y/2}(y)|.$$

Applying Vitali's covering lemma to $(B_{5s_y}(y))_{y \in F_2}$, we extract a subcovering $(B_{5s_j}(y_j))$ of F_2 such that the balls $B_{s_j}(y_j)$ are disjoint. For each j , define

$$I_j = \{i : x_i \in B_{s_j}(y_j)\}.$$

In particular, the sets I_j are disjoint. We claim that

$$(4.15) \quad s_j^\alpha \leq C_{N,\alpha} \sum_{i \in I_j} r_i^\alpha \quad \forall j \geq 1.$$

In order to establish (4.15), first observe that

$$(4.16) \quad A \cap B_{s_j/2}(y_j) \subset \bigcup_{i \in I_j} B_{r_i}(x_i).$$

In fact, given $z \in A \cap B_{s_j/2}(y_j)$, let i be such that $z \in B_{r_i}(x_i)$. We claim that $i \in I_j$. Assume, by way of contradiction, that $i \notin I_j$, i.e., suppose $x_i \notin B_{s_j}(y_j)$. Since

$$s_j \leq d(x_i, y_j) \leq d(x_i, z) + d(z, y) < r_i + s_j/2,$$

we have $s_j/2 < r_i$ and then $d(x_i, y_j) < 2r_i$. In other words, $y_j \in B_{2r_i}(x_i)$, which contradicts the definition of F_2 , since $y_j \in F_2$. This establishes (4.16). Applying (4.14) and (4.16), we have

$$\left(\frac{s_j}{2} \right)^N = \frac{1}{\omega_N} |B_{s_j/2}(y_j)| \leq \frac{8}{\omega_N} |A \cap B_{s_j/2}(y_j)| \leq \frac{8}{\omega_N} \sum_{i \in I_j} |B_{r_i}(x_i)| = 8 \sum_{i \in I_j} r_i^N.$$

Since $0 < \alpha \leq N$, we conclude that (4.15) holds. It now follows from (4.15) that

$$(4.17) \quad \mathcal{H}_\infty^\alpha(F_2) \leq \sum_{j=1}^{\infty} (5s_j)^\alpha \leq 5^\alpha C_{N,\alpha} \sum_{j=1}^{\infty} \sum_{i \in I_j} r_i^\alpha \leq C \sum_i r_i^\alpha \leq C \left[\mathcal{H}_\infty^\alpha(A) + \varepsilon \right].$$

Combining (4.13) and (4.17), we obtain

$$\mathcal{H}_\infty^\alpha(F) \leq C \left[\mathcal{H}_\infty^\alpha(A) + \varepsilon \right].$$

Since $\varepsilon > 0$ was arbitrary, (4.11) follows. \square

We now present the

Proof of Theorem 12 when $0 < \beta < 1$. Without loss of generality, we may assume that $v = 0$ on $\partial\Omega$; the general case follows by taking $v\varphi$, where φ is any function such that $\varphi \in C_c^\infty(\Omega)$ and $0 \leq \varphi \leq 1$ in Ω .

Fix $k \geq 1$ and let A_k be the set given by Lemma 3. We have

$$[v = 1] \subset A_k \cup E_k,$$

where $E_k = [v = 1] \setminus A_k$. We further decompose E_k as

$$E_k = E_{k,1} \cup E_{k,2},$$

where

$$E_{k,1} = \{x \in E_k : \Theta^*(x, A_k) \geq 1/4\} \quad \text{and} \quad E_{k,2} = \{x \in E_k : \Theta^*(x, A_k) < 1/4\}.$$

By Lemma 4, we have

$$(4.18) \quad \mathcal{H}_\infty^{N-2+\beta}(E_{k,1}) \leq C \mathcal{H}_\infty^{N-2+\beta}(A_k).$$

We now claim that

$$(4.19) \quad \limsup_{t \rightarrow 0} \frac{1}{t^{N-2+\beta}} \int_{B_t(x)} g(v) > 0 \quad \forall x \in E_{k,2}.$$

In fact, given $x \in E_{k,2}$, let $t_0 > 0$ be so small that

$$(4.20) \quad |A_k \cap B_t(x)| \leq \frac{1}{4} |B_t(x)|, \quad t \in (0, t_0).$$

Recall that $x \in \Omega \setminus A_k$ and $v(x) = 1$. It follows from (4.3) that

$$v(y) \geq 1 - Ck |x - y|^\beta \geq 1 - Ck t^\beta \quad \forall y \in B_t(x) \setminus A_k.$$

Since (1.4) holds, there exist $\tilde{C}_k > 0$ and a sequence $t_n \downarrow 0$ such that

$$g(1 - Ck t_n^\beta) \geq \frac{\tilde{C}_k}{t_n^{2-\beta}}, \quad \forall n \geq 1.$$

Thus, for all $n \geq 1$ sufficiently large, we get

$$g(v(y)) \geq \frac{\tilde{C}_k}{t_n^{2-\beta}} \quad \forall y \in B_{t_n}(x) \setminus A_k.$$

Since (4.20) holds, we obtain

$$\int_{B_{t_n}(x)} g(v) \geq \int_{B_{t_n}(x) \setminus A_k} g(v) \geq \frac{3}{4} |B_{t_n}(x)| \frac{\tilde{C}_k}{t_n^{2-\beta}} = C_k t_n^{N-2+\beta},$$

which gives (4.19). It now follows from (4.19) that $\mathcal{H}^{N-2+\beta}(E_{k,2}) = 0$ (see, e.g., [15, p. 77]); equivalently, we have

$$(4.21) \quad \mathcal{H}_\infty^{N-2+\beta}(E_{k,2}) = 0.$$

We now deduce from (4.18) and (4.21) that

$$\mathcal{H}_\infty^{N-2+\beta}(E_k) \leq C \mathcal{H}_\infty^{N-2+\beta}(A_k).$$

Therefore,

$$\mathcal{H}_\infty^{N-2+\beta}([v = 1]) \leq C \mathcal{H}_\infty^{N-2+\beta}(A_k) \leq \frac{C}{k} \|\nu\|_{\mathcal{M}}.$$

Since this estimate holds for every $k \geq 1$, as we let $k \rightarrow +\infty$ we obtain

$$\mathcal{H}_\infty^{N-2+\beta}([v = 1]) = 0.$$

In view of (4.5), the result follows. \square

The proof of Theorem 12 in the case $1 \leq \beta < 2$ follows the same strategy but is more technical. For this reason, we indicate the main steps in the proof. The counterpart of Lemma 3 is given by

Lemma 5. *Let $\nu \in \mathcal{M}(\Omega)$ and let v be the solution of (4.2). Given $1 \leq \beta < 2$ and $k \geq 1$, there exists a Borel set $A_k \subset \Omega$ such that*

$$(4.22) \quad \left| 2v\left(\frac{x+y}{2}\right) - v(x) - v(y) \right| \leq Ck|x-y|^\beta$$

for every $x, y \in \Omega \setminus A_k$ such that $(x+y)/2 \in \Omega \setminus A_k$; moreover,

$$(4.23) \quad \mathcal{H}_\infty^{N-2+\beta}(A_k) \leq \frac{C}{k} \|\nu\|_{\mathcal{M}},$$

for some constant $C > 0$ independent of k .

Proof. It suffices to consider the case where $\nu \geq 0$. Let A_k be given by (4.6). Proceeding as in the proof of Lemma 3, we obtain (4.23). Assume $N \geq 3$. We now show that w defined by

$$w(x) = a_N \int_\Omega \frac{d\nu(z)}{|z-x|^{N-2}} \quad \forall x \in \Omega,$$

where $a_N = 1/(N(N-2)\omega_N)$, satisfies property (4.22). Let $x, y \in \Omega \setminus A_k$ be such that $(x+y)/2 \in \Omega \setminus A_k$. Set $\delta = |x-y|$. We have

$$\left| 2w\left(\frac{x+y}{2}\right) - w(x) - w(y) \right| \leq a_N \int_\Omega \left| \frac{2}{\left|z - \frac{x+y}{2}\right|^{N-2}} - \frac{1}{|z-x|^{N-2}} - \frac{1}{|z-y|^{N-2}} \right| d\mu(z).$$

We split this integral into two parts:

$$\frac{1}{a_N} \left| 2w\left(\frac{x+y}{2}\right) - w(x) - w(y) \right| \leq \int_{|z-(x+y)/2| < 2\delta} + \int_{|z-(x+y)/2| \geq 2\delta}.$$

Note that $B_{2\delta}\left(\frac{x+y}{2}\right) \subset B_{5\delta/2}(x) \cap B_{5\delta/2}(y)$. Thus,

$$\begin{aligned} \int_{|z-(x+y)/2| < 2\delta} &\leq 2 \int_{B_{2\delta}\left(\frac{x+y}{2}\right)} \frac{d\nu(z)}{\left|z - \frac{x+y}{2}\right|^{N-2}} + \int_{B_{5\delta/2}(x)} \frac{d\nu(z)}{|z-x|^{N-2}} + \int_{B_{5\delta/2}(y)} \frac{d\nu(z)}{|z-y|^{N-2}} \\ &\leq C \int_0^{5\delta/2} \left[2\nu(B_s\left(\frac{x+y}{2}\right)) + \nu(B_s(x)) + \nu(B_s(y)) \right] \frac{ds}{s^{N-1}} \\ &\leq Ck\delta^\beta. \end{aligned}$$

On the other hand, we have

$$\left| \frac{2}{\left|z - \frac{x+y}{2}\right|^{N-2}} - \frac{1}{|z-x|^{N-2}} - \frac{1}{|z-y|^{N-2}} \right| \leq C \frac{\delta^2}{\left|z - \frac{x+y}{2}\right|^N}$$

if $|z - (x+y)/2| \geq 2\delta$. Therefore,

$$\begin{aligned} \int_{|z-(x+y)/2| \geq 2\delta} &\leq C\delta^2 \int_{|z-(x+y)/2| \geq 2\delta} \frac{d\nu(z)}{\left|z - \frac{x+y}{2}\right|^N} \\ &\leq CN\delta^2 \int_{2\delta}^\infty \nu(B_s\left(\frac{x+y}{2}\right)) \frac{ds}{s^{N+1}} \\ &\leq CNk\delta^2 \int_{2\delta}^\infty \frac{ds}{s^{3-\beta}} \leq Ck\delta^\beta. \end{aligned}$$

As in the proof of Lemma 3, we conclude that (4.22) holds. \square

We now present the

Proof of Theorem 12 completed. Assume $1 \leq \beta < 2$. Let $E_{k,1}$ and $E_{k,2}$ be defined as in the case $0 < \beta < 1$; in particular,

$$[v = 1] \subset A_k \cup E_{k,1} \cup E_{k,2}.$$

By Lemma 4, we have

$$\mathcal{H}_\infty^{N-2+\beta}(E_{k,1}) \leq C \mathcal{H}_\infty^{N-2+\beta}(A_k).$$

In order to establish the theorem, it suffices to prove (4.21). Given $x \in E_{2,k}$, let R_x denote the reflexion with respect to x , namely,

$$R_x(y) = 2x - y \quad \forall y \in \mathbb{R}^N.$$

We claim that

$$(4.24) \quad v(y) \geq 1 - Ck t^\beta \quad \forall y \in B_t(x) \setminus (A_k \cup R_x A_k).$$

In fact, for every $y \in B_t(x) \setminus (A_k \cup R_x A_k)$, we have $R_x(y) \in \Omega \setminus A_k$. Since $x \in \Omega \setminus A_k$, $v(x) = 1$ and $v \leq 1$, we get

$$v(y) \geq v(y) + v(R_x y) - 1 \geq 1 - Ck |x - y|^\beta \geq 1 - Ck t^\beta,$$

which is precisely (4.24). We now take $t_0 > 0$ so small that

$$|A_k \cap B_t(x)| \leq \frac{1}{4} |B_t(x)| \quad \forall t \in (0, t_0).$$

Therefore,

$$|(A_k \cup R_x A_k) \cap B_t(x)| \leq \frac{1}{2} |B_t(x)| \quad \forall t \in (0, t_0).$$

We can now proceed as in the case $0 < \beta < 1$ to conclude that

$$\limsup_{t \rightarrow 0} \frac{1}{t^{N-2+\beta}} \int_{B_t(x)} g(v) > 0 \quad \forall x \in E_{k,2}.$$

Thus,

$$\mathcal{H}_\infty^{N-2+\beta}(E_{k,2}) = 0.$$

As before, we deduce that (4.1) holds. The proof of Theorem 12 is complete. \square

5 Capacitary estimates related to problem (1.1)

In this section, we prove some estimates on the capacity of the set $[u^* = 1]$. They should be compared with the result of Theorem 12 concerning the Hausdorff measure of this set. Throughout this section, we assume that g satisfies a slightly stronger hypothesis than (1.4), namely

$$(5.1) \quad \liminf_{t \uparrow 1} \left\{ (1-t)^{(2-\beta)/\beta} g(t) \right\} > 0.$$

Given $p > 1$ and a Borel set $E \subset \Omega$, we denote by $\text{cap}_p(E)$ the capacity of E associated to $W_0^{1,p}(\Omega)$. Note that cap_2 coincides with the H^1 -capacity, denoted by cap elsewhere in this paper.

Our goal in this section is to establish

Theorem 13. *Let $v \in L^1(\Omega)$, $v \leq 1$ a.e., be such that $\Delta v \in \mathcal{M}(\Omega)$. Assume that g satisfies (5.1) for some $\beta \in (0, 1]$. If $g(v) \in L^1(\Omega)$, then*

$$(5.2) \quad \text{cap}_{2-\beta}([v = 1]) = 0.$$

Note that $\beta \in (0, 1]$ implies $2 - \beta \geq 1$, so that $\text{cap}_{2-\beta}$ is well-defined.

Proof.

Step 1. Proof of (5.2) if $v \in W_0^{1,1}(\Omega)$.

Set $\eta(s) = s^+ / (1 - s)$ and let $T_k(s) = \min\{s, k\}$. Since $v \in W_0^{1,1}(\Omega)$, for every $k \geq 1$, we have

$$(5.3) \quad \int_{\Omega} \nabla v \cdot \nabla T_k(\eta(v)) \, dx \leq k \|\Delta v\|_{\mathcal{M}}.$$

Indeed, inequality (5.3) (which formally amounts to multiplying Δv by $T_k(\eta(v))$) can be obtained by approximating v (e.g., through convolution) with smooth functions v_n such that $\|\Delta v_n\|_{\mathcal{M}} \leq \|\Delta v\|_{\mathcal{M}}$.

We can rewrite (5.3) as

$$(5.4) \quad \int_{[\eta(v) < k]} \frac{|\nabla v^+|^2}{(1-v)^2} \, dx \leq k \|\Delta v\|_{\mathcal{M}}.$$

On the other hand, applying Hölder's inequality with exponents $2/(2-\beta)$ and $2/\beta$, we have

$$(5.5) \quad \int_{[\eta(v) < k]} \frac{|\nabla v^+|^{2-\beta}}{(1-v)^{2(2-\beta)}} \, dx \leq \left(\int_{[\eta(v) < k]} \frac{|\nabla v^+|^2}{(1-v)^2} \, dx \right)^{1-\beta/2} \left(\int_{[\eta(v) < k]} \frac{1}{(1-v)^{2(2-\beta)/\beta}} \, dx \right)^{\beta/2}.$$

It then follows from (5.4)–(5.5) and the definition of η that

$$(5.6) \quad \int_{[\eta(v) < k]} \frac{|\nabla v^+|^{2-\beta}}{(1-v)^{2(2-\beta)}} \, dx \leq c(k \|\Delta v\|_{\mathcal{M}})^{1-\beta/2} \left(\int_{[\eta(v) < k]} 1 + \frac{\eta(v)^{(2-\beta)/\beta}}{(1-v)^{(2-\beta)/\beta}} \, dx \right)^{\beta/2}.$$

By assumption (5.1), there exists a constant $c_0 > 0$ such that

$$g(t)(1-t)^{(2-\beta)/\beta} \geq c_0 \quad \forall t \in (\frac{1}{2}, 1).$$

From (5.6), we obtain

$$\begin{aligned}
 (5.7) \quad & \int_{[\eta(v) < k]} \frac{|\nabla v^+|^{2-\beta}}{(1-v)^{2(2-\beta)}} dx \\
 & \leq c(k \|\Delta v\|_{\mathcal{M}})^{1-\beta/2} \left(\int_{[\eta(v) < k]} [1 + g(v) \eta(v)^{(2-\beta)/\beta}] dx \right)^{\beta/2} \\
 & \leq ck^{2-\beta} \|\Delta v\|_{\mathcal{M}}^{1-\beta/2} \left(\int_{\Omega} \frac{1 + g(v) T_k(\eta(v))^{(2-\beta)/\beta}}{k^{(2-\beta)/\beta}} dx \right)^{\beta/2}.
 \end{aligned}$$

Since $g(v) \in L^1(\Omega)$ (which also implies that $\eta(v)$ is finite a.e.), we have

$$\lim_{k \rightarrow +\infty} \int_{\Omega} \frac{1 + g(v) T_k(\eta(v))^{(2-\beta)/\beta}}{k^{(2-\beta)/\beta}} dx = 0.$$

It then follows from (5.7) that

$$(5.8) \quad \lim_{k \rightarrow +\infty} \int_{\Omega} \left| \frac{\nabla T_k(\eta(v))}{k} \right|^{2-\beta} dx = 0.$$

Note that

$$\frac{T_k(\eta(v))}{k} \geq 1 \quad \text{in } [\eta(v) \geq k].$$

Therefore,

$$\text{cap}_{2-\beta}([\eta(v) \geq k]) \leq \int_{\Omega} \left| \frac{\nabla T_k(\eta(v))}{k} \right|^{2-\beta} dx \xrightarrow{k \rightarrow +\infty} 0.$$

Since

$$[v = 1] = [\eta(v) = +\infty] = \bigcap_{k=1}^{\infty} [\eta(v) \geq k],$$

we conclude that

$$\text{cap}_{2-\beta}([v = 1]) = 0.$$

Step 2. Proof of Theorem 13 completed.

We replace v with $v\varphi$, where $\varphi \in C_c^\infty(\Omega)$ is a cut-off function, i.e., $0 \leq \varphi \leq 1$ in Ω and $\varphi = 1$ on a compact set $K \subset \Omega$. Since v and $\nabla v \in L^1_{\text{loc}}(\Omega)$, it follows that $\Delta(v\varphi) \in \mathcal{M}(\Omega)$. Moreover, $g(v\varphi) \leq g(v)$ a.e.; hence $g(v\varphi) \in L^1(\Omega)$. We can then apply the previous step to $v\varphi$ to deduce that $\text{cap}_{2-\beta}([v\varphi = 1]) = 0$. Thus

$$\text{cap}_{2-\beta}([v = 1] \cap K) = 0 \quad \text{for every compact } K \subset \Omega.$$

By the subadditivity of $\text{cap}_{2-\beta}$, we conclude that

$$\text{cap}_{2-\beta}([v = 1]) = 0. \quad \square$$

It is well-known (see, e.g., [15]) that $\text{cap}_1(E) = 0$ if and only if $\mathcal{H}^{N-1}(E) = 0$. Thus, in the case $\beta = 1$, we recover Theorem 12, but with a totally different proof. On the other hand, for any $p > 1$, $\text{cap}_p(E) = 0$ implies $\mathcal{H}^s(E) = 0$ for any $s > N - p$ (but the converse is not true). Thus, for $\beta \in (0, 1)$, Theorem 13 only gives $\mathcal{H}^s([v = 1]) = 0$ for any $s > N - 2 + \beta$, which is not optimal in view of Theorem 12.

However, it should be noticed that the proof of Theorem 13 only relies on energy estimates, which remain true for more general operators, for instance in the inhomogeneous case. Specifically, assume $A(x) = (a_{i,j}(x))$ is an $N \times N$ -matrix with bounded measurable coefficients satisfying

$$\lambda_1 |\xi|^2 \leq A(x)\xi \cdot \xi \leq \lambda_2 |\xi|^2 \quad \forall \xi \in \mathbb{R}^N, \quad \text{for a.e. } x \in \Omega,$$

where $0 < \lambda_1 \leq \lambda_2$. Proceeding as in the proof of Theorem 13, one obtains the following result.

Theorem 14. *Let $v \in L^1(\Omega)$ be such that $\text{div}(A(x)\nabla v)$ is a bounded measure “in the sense of Stampacchia”, i.e., assume there exists $\mu \in \mathcal{M}(\Omega)$ such that*

$$(5.9) \quad - \int_{\Omega} v \text{div}(A^*(x)\nabla \zeta) dx = \int_{\Omega} \zeta d\mu$$

for every $\zeta \in C_0(\overline{\Omega}) \cap H_0^1$ such that $\text{div}(A^*(x)\nabla \zeta) \in L^\infty(\Omega)$. Assume g satisfies (5.1) for some $\beta \in (0, 1]$. If $g(v) \in L^1(\Omega)$, then we have

$$\text{cap}_{2-\beta}([v = 1]) = 0.$$

Proof. Let μ_n be a suitable smooth convolution of μ , and consider the solutions v_n of

$$\begin{cases} -\text{div}(A(x)\nabla v_n) = \mu_n & \text{in } \Omega, \\ v_n \in H_0^1(\Omega). \end{cases}$$

Multiplying this equation by $T_k(\eta(v_n))$ (see the definition of $\eta(s)$ in Step 1 of Theorem 13), we get

$$\lambda_1 \int_{[\eta(v_n) < k]} \frac{|\nabla v_n^+|^2}{(1 - v_n)^2} dx \leq \int_{\Omega} (A(x)\nabla v_n) \cdot \nabla T_k(\eta(v_n)) dx \leq k \|\mu_n\|_{\mathcal{M}} \leq k \|\mu\|_{\mathcal{M}}.$$

Since the solutions in the sense of Stampacchia are unique and stable, v_n converges to v in $L^1(\Omega)$. Therefore, as $n \rightarrow +\infty$, we obtain

$$\lambda_1 \int_{[\eta(v) < k]} \frac{|\nabla v^+|^2}{(1-v)^2} dx \leq k \|\mu\|_{\mathcal{M}}.$$

One can now follow the argument in Step 1 of Theorem 13 in order to complete the proof. \square

In particular, if v satisfies the assumptions of Theorem 14, then

$$(5.10) \quad \mathcal{H}^s([v = 1]) = 0 \quad \text{for any } s > N - 2 + \beta, \quad \text{if } \beta \in (0, 1).$$

Note that

$$(5.11) \quad \mathcal{H}^{N-1}([v = 1]) = 0 \quad \text{if } \beta = 1.$$

It is an open problem whether (5.10) holds with $s = N - 2 + \beta$, where $\beta \in (0, 2)$, $\beta \neq 1$. Note that in the inhomogeneous case, it is not clear how to implement an approach based on Hölder continuity, as used in the proof of Theorem 12.

Remark 1. In the same spirit, the proof of Theorem 13 extends to nonlinear operators such as the p -Laplacian, for functions v which satisfy $-\operatorname{div}(|\nabla v|^{p-2} \nabla v) \in \mathcal{M}(\Omega)$ “in the renormalized sense” (see [9] for the precise definition). In this case, one can prove with the same method that if (5.1) holds for some $\beta \in (0, 1]$ and $g(v) \in L^1(\Omega)$, then

$$\operatorname{cap}_q([v = 1]) = 0 \quad \text{with } q = \frac{(2 - \beta)p}{2(1 - \beta) + \beta p}.$$

Note that if $\beta = 1$, one still has

$$\operatorname{cap}_1([v = 1]) = 0 = \mathcal{H}^{N-1}([v = 1]).$$

6 Every diffuse measure is good for some g

Our goal in this section is to establish the following theorem.

Theorem 15. *Let $\mu \in \mathcal{M}(\Omega)$ be such that μ^+ is diffuse. Then, there exists some g such that $\mu \in \mathcal{G}(g)$.*

We start with the

Proposition 5. *If $g_1 \leq g_2$, then $\mathcal{G}(g_1) \subset \mathcal{G}(g_2)$.*

Proof. Given $\mu \in \mathcal{G}(g_1)$, let u be the solution of

$$\begin{cases} -\Delta u + g_1(u) = \mu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Let μ^* be the reduced measure relative to g_2 and denote by u^* the solution of

$$\begin{cases} -\Delta u^* + g_2(u^*) = \mu^* & \text{in } \Omega, \\ u^* = 0 & \text{on } \partial\Omega. \end{cases}$$

Since $\mu^* \leq \mu$ and $g_2 \geq g_1$, we have $u^* \leq u$ (see [4, Corollary B.2]). In other words, $u - u^* \geq 0$ in Ω and $u - u^* = 0$ on the set $[u^* = 1]$. Thus, by (2.3), we have

$$(\mu^* - \mu)_d = [\Delta(u - u^*)]_d \geq 0 \quad \text{in } [u^* = 1].$$

This implies $(\mu^*)_d = \mu_d$ in $[u^* = 1]$. On the other hand, by Theorem 4,

$$(\mu^*)_d = \mu_d \quad \text{in } [u^* < 1].$$

We conclude that

$$(6.1) \quad (\mu^*)_d = \mu_d.$$

Finally, since μ is a good measure relative to g_1 , we have $\mu_c \leq 0$. Thus, by (2.10),

$$(6.2) \quad (\mu^*)_c = -(\mu_c)^- = \mu_c.$$

It follows from (6.1) and (6.2) that $\mu = \mu^* \in \mathcal{G}(g_2)$. This concludes the proof of the proposition. \square

Related to the previous result, we point out the following

Open Problem. Assume $g_1 \leq g_2$ and $\mathcal{G}(g_1) = \mathcal{G}(g_2)$. Is it true that $g_1 = g_2$?

Lemma 6. *Let $\mu \in \mathcal{M}(\Omega)$ be a nonnegative diffuse measure. Given $\varepsilon > 0$, $s_0 \in (0, 1)$, and a continuous nondecreasing function $g : (-\infty, 1) \rightarrow \mathbb{R}$ satisfying (1.2)–(1.3), there exists $\tilde{g} : (-\infty, 1) \rightarrow \mathbb{R}$ with*

$$(6.3) \quad \tilde{g} \geq g \quad \text{in } (-\infty, 1), \quad \tilde{g} = g \quad \text{in } (-\infty, s_0],$$

such that

$$(6.4) \quad \mu([v = 1]) < \varepsilon,$$

where v is the largest subsolution of the problem

$$\begin{cases} -\Delta u + \tilde{g}(u) = \mu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Proof. Fix $t_0 \in (s_0, 1)$. Let $g_k : (-\infty, 1) \rightarrow \mathbb{R}$ be any increasing sequence of continuous nondecreasing functions such that

$$\begin{cases} g_k(t) = g(t) & \text{if } t \leq s_0, \\ g_k(t) \geq k & \text{if } t \geq t_0, \\ g_k(t) \geq g(t) & \forall t \in (-\infty, 1). \end{cases}$$

For each $k \geq 1$, let μ_k denote the reduced measure of μ relative to g_k . We denote by v_k the corresponding solution. In particular, by Proposition 1,

$$(6.5) \quad \int_{\Omega} |\Delta v_k| \leq 2\|\mu_k\|_{\mathcal{M}} \leq 2\|\mu\|_{\mathcal{M}}$$

and

$$(6.6) \quad \int_{\Omega} g_k(v_k) \leq \|\mu\|_{\mathcal{M}}.$$

In view of (6.6), we have

$$(6.7) \quad |[v_k \geq t_0]| \leq \frac{1}{k} \int_{\Omega} g_k(v_k) \leq \frac{1}{k} \|\mu\|_{\mathcal{M}} \rightarrow 0$$

as $k \rightarrow +\infty$. On the other hand, the sequence (v_k) is non-increasing; thus, there exists $v \in L^1(\Omega)$ such that $v_k \downarrow v$ in $L^1(\Omega)$. By (6.7), we have $v \leq t_0$ a.e. Moreover, since $0 \leq v_k \leq 1$ a.e., it follows from (6.5) that (v_k) is bounded in $L^\infty(\Omega) \cap H_0^1(\Omega)$. We then conclude that $v_k \rightarrow v$ μ -a.e. in Ω (see, e.g., [6, Lemma 2.1]). Therefore,

$$\mu([v_k > t_0]) \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

The lemma then follows by taking $\tilde{g} = g_{k_0}$ for some $k_0 \geq 1$ sufficiently large. \square

We now present the

Proof of Theorem 15. We split the proof of the theorem into two steps.

Step 1. Given $\mu \in \mathcal{M}(\Omega)$ diffuse and nonnegative, there exists g satisfying (1.2)–(1.3) such that $\mu \in \mathcal{G}(g)$.

We begin by constructing a sequence (g_k) as follows. Let $g_0(t) = t/(1-t)$, $t \in [0, 1)$. Given g_k , we apply Lemma 6 to $g = g_k$, $\varepsilon = 1/2^k$ and $s_0 = 1 - 1/2^k$. Set $g_{k+1} = \tilde{g}$, where \tilde{g} is the function given by Lemma 6. Then the sequence (g_k) is nondecreasing and

$$g_k = g_{k_0} \quad \text{in } (-\infty, 1 - 1/2^{k_0}], \quad k \geq k_0.$$

Set

$$g(t) = \lim_{k \rightarrow +\infty} g_k(t), \quad t \in (-\infty, 1).$$

We claim that $\mu \in \mathcal{G}(g)$. Indeed, let μ_k denote the reduced measure of μ relative to g_k . Then μ_k is also a diffuse measure. Since $g \geq g_k$, it follows from Proposition 5 that $\mu_k \in \mathcal{G}(g)$ for every $k \geq 1$. Let v_k be the solution of

$$\begin{cases} -\Delta v_k + g_k(v_k) = \mu_k & \text{in } \Omega, \\ v_k = 0 & \text{on } \partial\Omega. \end{cases}$$

By Theorem 4 and the choice of g_k , we have

$$\|\mu - \mu_k\|_{\mathcal{M}} \leq \mu([v_k = 1]) \leq 1/2^k.$$

Thus,

$$\mu_k \rightarrow \mu \quad \text{strongly in } \mathcal{M}(\Omega).$$

Since $\mathcal{G}(g)$ is closed, we conclude that $\mu \in \mathcal{G}(g)$, as claimed.

Step 2. Proof of the theorem completed.

Let $\mu \in \mathcal{M}(\Omega)$ be such that $\mu_c \leq 0$; in other words, μ^+ is diffuse. We can then apply the previous step to μ^+ to conclude that there exists g such that $\mu^+ \in \mathcal{G}(g)$. Since $\mu \leq \mu^+$, it follows from Proposition 4 that μ is also good for g . \square

7 Measures which are good for every g

In this section, we characterize the set of measures which are always good. In order to do so, we first need to recall some notions about obstacle problems with measure data. Throughout this section, we denote by β any maximal monotone graph (m.m.g.) of the form

$$(7.1) \quad \beta(t) = \begin{cases} b(t) & \text{if } t < 1, \\ [b(1), \infty) & \text{if } t = 1, \\ \emptyset & \text{if } t > 1, \end{cases}$$

where $b : (-\infty, 1] \rightarrow \mathbb{R}$ is a nondecreasing continuous function such that $b(t) = 0$ if $t \leq 0$. Given a bounded measure μ in Ω , we say that w is a solution of

$$(7.2) \quad \begin{cases} -\Delta w + \beta(w) \ni \mu & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega, \end{cases}$$

if $w \in L^1(\Omega)$, $w \leq 1$ a.e., $\Delta w \in \mathcal{M}(\Omega)$, and there exists a nonnegative diffuse measure $\nu \in \mathcal{M}(\Omega)$ such that $\nu_a \in \beta(w)$ a.e., ν_s is concentrated on the set $[w = 1]$, and

$$(7.3) \quad - \int_{\Omega} w \Delta \zeta + \int_{\Omega} \zeta d\nu = \int_{\Omega} \zeta d\mu \quad \forall \zeta \in C_0^2(\overline{\Omega}).$$

(Here, ν_a and ν_s denote the absolutely continuous and singular parts of ν with respect to Lebesgue measure in \mathbb{R}^N .)

In particular, the measure $\mu + \Delta w$ is diffuse and

$$(7.4) \quad \mu + \Delta w = \nu \geq \inf \beta(1) \quad \text{in } [w = 1].$$

Problem (7.2) has been studied by Dall’Aglione–Leone [11], Dall’Aglione–Dal Maso [10], Brezis–Ponce [6]; see also the references therein. It turns out that (7.2) has a solution if and only if μ^+ is diffuse; moreover, this solution is unique and is the largest solution of the problem

$$(7.5) \quad \begin{cases} -\Delta v + b(v) \leq \mu & \text{in } \Omega, \\ v \leq 1 & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Our goal in this section is to establish the following.

Theorem 16. *Let $\mu \in \mathcal{M}(\Omega)$. Then μ is good for every g if and only if μ^+ is diffuse and*

$$\mu + \Delta w_0 \in L^1(\Omega),$$

where w_0 is the unique solution of the obstacle problem

$$(7.6) \quad \begin{cases} -\Delta w_0 + \beta_0(w_0) \ni \mu & \text{in } \Omega, \\ w_0 = 0 & \text{on } \partial\Omega, \end{cases}$$

with $\beta_0(s) = 0$ if $s < 1$ and $\beta_0(1) = [0, \infty)$.

Author query:
This remark does not appear in the hard copy you submitted, do you want to leave it?

Remark 2. It is known from [2] that if $\mu \in L^1(\Omega)$, then problem (1.1) has a solution for every g . This is consistent with Theorem 16. Indeed, let w_0 be the solution of (7.6) with $\mu \in L^1(\Omega)$. Then, in view of (2.3), we have

$$\mu + \Delta w_0 \leq \mu \quad \text{on } [w_0 = 1].$$

Since $\mu + \Delta w_0$ is a nonnegative measure and it is concentrated on $[w_0 = 1]$, we conclude that

$$0 \leq \mu + \Delta w_0 \leq \mu \quad \text{in } \Omega.$$

Hence, $\mu \in L^1(\Omega)$ implies that $\mu + \Delta w_0 \in L^1(\Omega)$.

For the proof of Theorem 16, we require the following lemmas.

Lemma 7. *Let β be a m.m.g. and let $\mu \in \mathcal{M}(\Omega)$ be such that μ^+ is diffuse. If*

$$(7.7) \quad \mu_a^+ \in L^\infty(\Omega) \quad \text{and} \quad \|\mu_a^+\|_{L^\infty} < \inf \beta(1),$$

then

$$(7.8) \quad |[w = 1]| = 0,$$

where w is the solution of (7.2).

Proof. By (2.3), we have $(\Delta w)_d \leq 0$ on the set $[w = 1]$. Thus,

$$\mu_d \geq (\mu + \Delta w)_d = \mu + \Delta w \geq \inf \beta(1) \quad \text{in } [w = 1].$$

Comparing the absolutely continuous part of both sides, we get

$$\mu_a = (\mu_d)_a \geq \inf \beta(1) \quad \text{a.e. in } [w = 1].$$

In view of (7.7), we deduce that (7.8) holds. \square

Lemma 8. *Let $\mu \in \mathcal{M}(\Omega)$ be such that μ^+ is diffuse. Given two m.m.g. β_1, β_2 , let w_i be the solution of (7.2) associated to β_i , $i = 1, 2$. If $\beta_1 \geq \beta_2$, then*

$$(7.9) \quad 0 \leq \mu + \Delta w_1 \leq \mu + \Delta w_2 \quad \text{in } [w_1 = 1].$$

Proof. By comparison, we have $w_2 - w_1 \geq 0$ a.e. In particular, $w_2 - w_1 = 0$ in $[w_1 = 1]$. Applying (2.3), we get

$$[\Delta(w_2 - w_1)]_d \geq 0 \quad \text{in } [w_1 = 1].$$

Thus, on the set $[w_1 = 1]$, we have

$$\mu + \Delta w_1 = (\mu + \Delta w_1)_d \leq (\mu + \Delta w_2)_d = \mu + \Delta w_2.$$

Since w_1 is the solution of (7.2) with $\beta = \beta_1$, we have $\nu_1 = \mu + \Delta w_1 \geq 0$ in Ω . We conclude that (7.9) holds. \square

Proof of Theorem 16.

Proof of (\Leftarrow) . We establish a slightly more general result:

Proposition 6. *Let $\mu \in \mathcal{M}(\Omega)$ be such that μ^+ is diffuse. Assume that*

$$(7.10) \quad \mu + \Delta w \in L^1(\Omega),$$

where w is the unique solution of (7.2). Then μ is good for every g such that $g \geq \beta$.

Proof. We first assume $\mu_a^+ \in L^\infty(\Omega)$. Let β_1 be a m.m.g. such that

$$\beta \leq \beta_1 \leq g \quad \text{and} \quad \|\mu_a^+\|_{L^\infty} < \inf \beta_1(1).$$

Let w_1 be the solution of (7.2) with obstacle β_1 . We claim that

$$(7.11) \quad \mu + \Delta w_1 \in L^1(\Omega).$$

In fact, since w_1 is the solution of an obstacle problem, the measure $(\mu + \Delta w_1)_s$ is concentrated on the set $[w_1 = 1]$. By Lemma 8 above, we have

$$0 \leq (\mu + \Delta w_1)_s \leq (\mu + \Delta w)_s = 0 \quad \text{in } [w_1 = 1].$$

This establishes (7.11).

On the other hand, it follows from Lemma 7 that the set $[w_1 = 1]$ has zero Lebesgue measure. We conclude that

$$\mu + \Delta w_1 = b_1(w_1) \quad \text{a.e.}$$

In other words, w_1 verifies

$$\begin{cases} -\Delta w_1 + b_1(w_1) = \mu & \text{in } \Omega, \\ w_1 = 0 & \text{on } \partial\Omega. \end{cases}$$

Since $|[w_1 = 1]| = 0$, by a variant of the De La Vallée Poussin theorem (see [12, Remark 23] or [13, Theorem II.22]), one can find g_1 satisfying (1.2) such that

$$b_1 \leq g_1 \leq g \quad \text{and} \quad g_1(w_1) \in L^1(\Omega).$$

Thus, in view of Proposition 4, μ is a good measure for g_1 . By Proposition 5, we conclude that μ is also good for g . Since $g \geq \beta$ was arbitrary, the result follows when $\mu_a^+ \in L^\infty(\Omega)$.

In order to establish the proposition for any measure μ satisfying (7.10), we let

$$\mu_n = \min \{\mu_a, n\} + \mu_s, \quad n \geq 1.$$

Proceeding as in the proof of (7.11), we have for every $n \geq 1$,

$$\mu_n + \Delta w_n \in L^1(\Omega),$$

where w_n is the solution of (7.2) with data μ_n . Moreover, $(\mu_n)_a^+ \in L^\infty(\Omega)$. Thus, μ_n is good for every $g \geq \beta$. Letting $n \rightarrow +\infty$, we conclude that μ is also good for any such g . \square

Proof of (\Rightarrow) . We require the following.

Lemma 9. *Let $\mu \in \mathcal{M}(\Omega)$. Assume that μ is good for every g , and that*

$$(7.12) \quad \mu_a^+ \in L^\infty(\Omega) \quad \text{and} \quad \|\mu_a^+\|_{L^\infty} \leq \inf \beta(1).$$

Let w be the solution of (7.2). Then

$$(7.13) \quad \begin{cases} -\Delta w + b(w) = \mu & \text{in } \Omega, \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

Proof. By making a small perturbation of μ , it suffices to establish the result when

$$(7.14) \quad \|\mu_a^+\|_{L^\infty} < \inf \beta(1).$$

By Lemma 7 above, $|[w = 1]| = 0$. Thus, one can find a continuous nondecreasing function $H : (-\infty, 1) \rightarrow \mathbb{R}$, $H \geq b$, satisfying (1.2) and such that $H(w) \in L^1(\Omega)$. We now take a sequence of functions (g_n) such that

$$g_n \leq H \quad \text{for all } n \geq 1 \quad \text{and} \quad g_n \downarrow b \quad \text{as } n \uparrow +\infty.$$

Since μ is good for every g_n , there exist v_n satisfying (1.1) with nonlinearity g_n . Clearly, $v_n \uparrow v$, where $v \in L^1(\Omega)$ and $v \leq 1$ a.e. By Fatou, v verifies (7.5); in particular, $v \leq w$ a.e. Thus

$$(7.15) \quad |[v = 1]| \leq |[w = 1]| = 0.$$

On the other hand,

$$(7.16) \quad g_n(v_n) \rightarrow b(v) \quad \text{a.e. on } [v < 1].$$

Thus, by (7.15)–(7.16), we have

$$g_n(v_n) \rightarrow b(v) \quad \text{a.e.}$$

Since $g_n(v_n) \leq H(w)$ a.e., it follows by dominated convergence that

$$g_n(v_n) \rightarrow b(v) \quad \text{in } L^1(\Omega).$$

We deduce that v satisfies (7.13). In particular, v is also a solution of (7.2). By uniqueness, we conclude that $v = w$. This establishes the lemma. \square

We can now conclude the proof of Theorem 16.

Let μ be a measure which is good for every g . We assume in addition that $\mu_a^+ \in L^\infty(\Omega)$. Let $b_n : (-\infty, 1] \rightarrow \mathbb{R}$ be a sequence of nondecreasing continuous functions such that $b_n(t) = 0$ if $t \leq 0$, $b_n(1) = \|\mu_a^+\|_{L^\infty}$ and $b_n(t) \downarrow 0$ uniformly away from $t = 1$. By Lemma 9, equation (7.13) has a solution $v_n \leq 1$ associated to b_n . Note that $v_n \uparrow v$, where $v \in L^1(\Omega)$, $v \leq 1$ a.e. Moreover, passing to a subsequence if necessary, we have

$$b_n(v_n) \rightharpoonup f \quad \text{weakly in } L^\infty(\Omega)$$

for some $f \in L^\infty(\Omega)$ with $\|f\|_{L^\infty} \leq \|\mu_a^+\|_{L^\infty}$.

We claim that $v = w_0$ a.e. Indeed, v satisfies

$$\begin{cases} -\Delta v = \mu - f & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Since

$$0 \leq b_n(v_n) \leq b_n(v) \quad \text{a.e.} \quad \forall n \geq 1,$$

letting $n \rightarrow +\infty$, we obtain

$$0 \leq f \leq \alpha \chi_{[v=1]} \quad \text{a.e.},$$

where $\alpha = \|\mu_a^+\|_{L^\infty}$. This implies that f is nonnegative and concentrated on the set $[v = 1]$. Therefore, v satisfies (7.6), so $v = w_0$ as claimed. We conclude that

$$\mu + \Delta w_0 = f \in L^\infty(\Omega).$$

This establishes the theorem under the additional assumption that $\mu_a^+ \in L^\infty(\Omega)$. The general case follows easily by an approximation argument. \square

Before proving Theorem 3, we start with the following

Proposition 7. *Given $\mu \in \mathcal{M}(\Omega)$, let v be the unique solution of*

$$(7.17) \quad \begin{cases} -\Delta v = \mu & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

If $v \leq 1$ a.e., then μ is good for every g .

Proof. Let $\alpha < 1$. Since $v \leq 1$ a.e., αv is a supersolution of problem (1.1) with data $\alpha\mu$. Thus, by Proposition 4, $\alpha\mu$ is good for every $\alpha < 1$. Since $\mathcal{G}(g)$ is closed with respect to the strong topology in $\mathcal{M}(\Omega)$, we conclude that $\mu \in \mathcal{G}(g)$ for every g . \square

We now present the

Proof of Theorem 3. The implication “ \Leftarrow ” already follows from Proposition 7 above. We now establish the reverse implication. Let μ be a singular measure which is good for every g . In particular, μ^+ is diffuse. Let $\nu = \mu + \Delta w_0$, where w_0 is the solution of (7.6). In view of the definition of β_0 , ν is a nonnegative diffuse measure concentrated on the set $[w_0 = 1]$. By (2.3), on this set we have $(\Delta w_0)_d \leq 0$, so that

$$0 \leq \nu \leq \mu.$$

On the other hand, by Theorem 16, $\nu \in L^1(\Omega)$. Since μ is singular, we conclude that $\nu = 0$; hence w_0 coincides with the unique solution v of (7.17). Since $w_0 \leq 1$ a.e., the result follows. \square

8 How to construct diffuse measures which are not good

Our goal in this section is to establish Theorem 1. The main ingredient in the proof is the following.

Lemma 10. *Given g , there exists $v \in C_0(\overline{\Omega})$ such that*

$$(8.1) \quad \Delta v \in L^1(\Omega), \quad v \leq 1 \text{ in } \Omega, \quad \text{cap}([v = 1]) > 0 \quad \text{and} \quad g(v) \in L^1(\Omega).$$

Proof. Let (ℓ_k) be a decreasing sequence of positive numbers such that

$$(8.2) \quad \ell_k \leq \theta \ell_{k-1} \quad \forall k \geq 2,$$

for some $\theta \in (0, \frac{1}{2})$. Let (k_j) be an increasing sequence of nonnegative integers. Both sequences (ℓ_k) and (k_j) will be explicitly chosen later on.

We now recall briefly the construction in [18] of the Cantor set F associated to the subsequence (ℓ_{k_j}) . We assume for simplicity that $\Omega = Q_1$, the unit cube centered at 0.

We first define a decreasing sequence of sets $(F_j)_{j \geq 0}$ as follows. Let $F_0 = Q_1$, $k_0 = 0$ and $\ell_0 = 1$. We now proceed by induction. Assume F_{j-1} , $j \geq 1$, is the union of $2^{N k_{j-1}}$ disjoint cubes of length $\ell_{k_{j-1}}$. Let Q_i be any component of F_{j-1} , and let $\tilde{Q}_i \subset Q_i$ be a smaller cube concentric to Q_i (so that the ratio between their lengths is $\frac{1}{2} + \theta \in (\frac{1}{2}, 1)$). Inside \tilde{Q}_i , we select $2^{N(k_j - k_{j-1})}$ cubes $Q_{i,s}$ of length ℓ_{k_j} , uniformly distributed in \tilde{Q}_i . Set

$$F_j = \bigcup_{i,s} Q_{i,s}.$$

Thus, $F_j \subset F_{j-1}$, and F_j is the union of 2^{Nk_j} disjoint cubes of length ℓ_{k_j} . The Cantor set associated to the subsequence (ℓ_{k_j}) is then defined as

$$F = \bigcap_{j=1}^{\infty} F_j.$$

We now split the proof of the lemma into two cases, $N \geq 3$ or $N = 2$.

Case 1. $N \geq 3$

We start with the following

Claim 1. *For every $j \geq 1$, we have*

$$(8.3) \quad \text{cap}(F_j, F_{j-1}) \leq C_\theta 2^{Nk_j} \ell_{k_j}^{N-2},$$

where $\text{cap}(F_j, F_{j-1})$ denotes the H^1 -capacity of the set F_j with respect to F_{j-1} .

Proof. Since F_{j-1} has $2^{Nk_{j-1}}$ connected components, it suffices to show that

$$(8.4) \quad \text{cap}(F_j \cap Q_i, Q_i) \leq C_\theta 2^{N(k_j - k_{j-1})} \ell_{k_j}^{N-2},$$

where Q_i is any component of F_{j-1} . Note that each component $Q_{i,s}$ of $F_j \cap Q_i$ has length ℓ_{k_j} and (see [18])

$$d(Q_{i,s}, \partial Q_i) \geq \frac{1-2\theta}{4} \text{diam } Q_i.$$

Hence

$$(8.5) \quad \text{cap}(Q_{i,s}, Q_i) \leq C_\theta \ell_{k_j}^{N-2}.$$

Recall that Q_i contains $2^{N(k_j - k_{j-1})}$ components $Q_{i,s}$. By the subadditivity of the capacity, we conclude that (8.4) holds. This concludes the proof of the claim. \square

By (8.3) and Theorem E.1 in [4], there exists $v_j \in C_c^\infty(F_{j-1})$ such that $0 \leq v_j \leq 1$ in Ω , $v_j = 1$ on F_j , and

$$(8.6) \quad \int_{\Omega} |\Delta v_j| \leq C 2^{Nk_j} \ell_{k_j}^{N-2}.$$

Our aim is to construct the function v of the form

$$(8.7) \quad v = \sum_{j=1}^{\infty} \alpha_j v_j,$$

where (α_j) is a sequence of positive numbers to be chosen later on such that

$$(8.8) \quad \sum_{j=1}^{\infty} \alpha_j = 1.$$

Clearly, $v \in C_0(\overline{\Omega})$ and $v \leq 1$ in Ω . Moreover, $v = 1$ precisely on $\bigcap_j F_j = F$.

We claim that one can choose (ℓ_k) , (k_j) and (α_j) such that

$$(8.9) \quad \sum_{j=1}^{\infty} \frac{1}{2^{Nk_j} \ell_{k_j}^{N-2}} < \infty,$$

$$(8.10) \quad \sum_{j=1}^{\infty} \alpha_j 2^{Nk_j} \ell_{k_j}^{N-2} < \infty,$$

$$(8.11) \quad \sum_{j=1}^{\infty} g \left(\sum_{i=1}^j \alpha_i \right) 2^{Nk_j} \ell_{k_j}^N < \infty.$$

In fact, let

$$\alpha_j = 3 \cdot 2^{-2j} \quad \forall j \geq 1,$$

so that (8.8) holds. Let (k_j) be any increasing sequence of positive numbers such that

$$\frac{g(1 - 2^{-2j})}{2^{\frac{N}{N-2}k_j}} \leq \frac{1}{2^j} \quad \forall j \geq 1.$$

Finally, we take (ℓ_k) satisfying (8.2) (with, say, $\theta = \frac{3}{4}$) and

$$2^{Nk_j} \ell_{k_j}^{N-2} = 2^j \quad \forall j \geq 1.$$

It is immediate that (8.9) and (8.10) hold. After some straightforward computation, the left-hand side of (8.11) can be estimated by

$$\sum_{j=1}^{\infty} g(1 - 2^{-2j}) \left(\frac{2^j}{2^{2k_j}} \right)^{\frac{N}{N-2}} \leq \sum_{j=1}^{\infty} \frac{g(1 - 2^{-2j})}{2^{\frac{N}{N-2}k_j}},$$

which is finite in view of our choice of (k_j) . We conclude that (8.9)–(8.11) hold.

By construction, $[v = 1]$ coincides with F . On the other hand, by [18], $\text{cap}(F) > 0$ if and only if

$$(8.12) \quad \sum_{j=1}^{\infty} \frac{1}{2^{Nk_j} \ell_{k_j}^{N-2}} < \infty.$$

Thus, in view of (8.9), $\text{cap}(F) > 0$, so

$$(8.13) \quad \text{cap}([v = 1]) > 0.$$

By (8.6), (8.7) and (8.10), we have

$$(8.14) \quad \Delta v \in L^1(\Omega).$$

Finally, it follows from (8.11) that

$$(8.15) \quad g(v) \in L^1(\Omega).$$

The proof of the lemma is complete when $N \geq 3$.

Case 2. $N = 2$.

We start with the

Claim 2. *For every $j \geq 1$, we have*

$$(8.16) \quad \text{cap}(F_j, F_{j-1}) \leq C_\theta 4^{k_j} \left(\log \frac{1}{\ell_{k_j}} \right)^{-1}.$$

The argument is similar to the proof of Claim 1. It suffices to observe that the analogue of (8.5) is

$$(8.17) \quad \text{cap}(Q_{i,s}, Q_i) \leq C_\theta \left(\log \frac{1}{\ell_{k_j}} \right)^{-1}.$$

We now conclude the proof of the lemma. Let (α_j) be defined as before. Take an increasing sequence of positive integers (k_j) such that

$$\frac{g(1 - 2^{-2j})}{4^{4^{k_j}}} \leq \frac{1}{2^j} \quad \forall j \geq 1.$$

Finally, let (ℓ_k) satisfy (8.2) and

$$\ell_{k_j} = 4^{-4^{k_j} \cdot 2^{-j}} \quad \forall j \geq 1.$$

With such choices, one can easily check that

$$(8.18) \quad \sum_{j=1}^{\infty} \frac{1}{4^{k_j}} \log \frac{1}{\ell_{k_j}} < \infty,$$

$$(8.19) \quad \sum_{j=1}^{\infty} \alpha_j 4^{k_j} \left(\log \frac{1}{\ell_{k_j}} \right)^{-1} < \infty,$$

$$(8.20) \quad \sum_{j=1}^{\infty} g \left(\sum_{i=1}^j \alpha_i \right) 4^{k_j} \ell_{k_j}^2 < \infty.$$

Let v be given by (8.7), where $v_j \in C_c^\infty(F_{j-1})$ is such that $0 \leq v_j \leq 1$ in Ω , $v_j = 1$ on F_j , and

$$\int_{\Omega} |\Delta v_j| \leq C 4^{k_j} \left(\log \frac{1}{\ell_{k_j}} \right)^{-1}.$$

(The existence of such v_j follows from Claim 2 above.) In particular, $[v = 1] = F$. By (8.18), we have (see [18, Lemma 4])

$$\text{cap}([v = 1]) > 0.$$

Moreover, proceeding as before, we deduce from (8.19) and (8.20) that

$$\Delta v \in L^1(\Omega) \quad \text{and} \quad g(v) \in L^1(\Omega).$$

This concludes the proof of the lemma. \square

Lemma 10 allows us to prove the following result.

Proposition 8. *For every g , there exist a nonnegative function $h_0 \in L^1(\Omega)$ and a compact set $K_0 \subset \Omega$, with $|K_0| = 0$ and $\text{cap}(K_0) > 0$, such that for any measure $\sigma \geq 0$ supported in K_0 , we have*

$$(8.21) \quad (h_0 + \sigma)^* = h_0,$$

where $(h_0 + \sigma)^*$ is the reduced measure associated to $h_0 + \sigma$.

Proof. Take $K_0 = [v = 1]$ and $f_0 = -\Delta v + g(v)$, where v is the function constructed in Lemma 10. We begin with the following

Claim. *If λ is a good measure $\geq f_0$, then $\lambda(K_0) = 0$.*

Proof. We first observe that λ is a diffuse measure. In fact, since λ is good, we have $\lambda_c \leq 0$ by Corollary 2. On the other hand, $\lambda \geq f_0$ implies $\lambda_c \geq 0$. Thus, $\lambda_c = 0$, so that λ is diffuse. Let u be the solution of

$$\begin{cases} -\Delta u + g(u) = \lambda & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Clearly, $u \geq v$ a.e. Moreover, since $v = 1$ in K_0 , we have $u - v = 0$ in K_0 . Thus, by (2.3),

$$f_0 - \lambda = \Delta(u - v) = [\Delta(u - v)]_d \geq 0 \quad \text{in } K_0.$$

In other words, $\lambda \leq f_0$ in K_0 ; thus, $\lambda = f_0$ in K_0 . Since $|K_0| = 0$, we deduce that $\lambda(K_0) = 0$. This establishes the claim. \square

Let $h_0 = f_0^+$. We now show that h_0 and K_0 satisfy the desired properties. In fact, let $\sigma \geq 0$ be a measure concentrated on K_0 . Since $h_0 + \sigma \geq 0$, it follows from Corollary 3 that

$$0 \leq (h_0 + \sigma)^* \leq h_0 + \sigma \quad \text{in } \Omega.$$

Moreover, by Corollary 1,

$$[(h_0 + \sigma)^*]_{\mathfrak{a}} = h_0 \quad \text{a.e. in } \Omega.$$

Thus

$$f_0 \leq h_0 \leq (h_0 + \sigma)^* \leq h_0 + \sigma \quad \text{in } \Omega.$$

In particular, $[(h_0 + \sigma)^*]_{\mathfrak{s}}$ is concentrated on K_0 . By our previous claim, we have $[(h_0 + \sigma)^*]_{\mathfrak{s}} = 0$ in K_0 . Therefore,

$$(h_0 + \sigma)^* = [(h_0 + \sigma)^*]_{\mathfrak{a}} = h_0 \quad \text{in } \Omega. \quad \square$$

Remark 3. A slight modification in the construction of v , given by Lemma 10, allows us to obtain the following additional property in the statement of Proposition 8: given any $\varepsilon > 0$ and any ball $B_r \subset\subset \Omega$, one can choose h_0 and K_0 such that

$$\|h_0\|_{L^1} < \varepsilon \quad \text{and} \quad K_0 \subset B_r.$$

Theorem 1 is now a consequence of Proposition 8.

Proof of Theorem 1. Given g , let h_0, K_0 be as in the statement of Proposition 8. Since $\text{cap}(K_0) > 0$, there exists a diffuse measure $\sigma \geq 0$ concentrated on K_0 such that $\sigma(K_0) = 1$ (see, e.g., [8]). Let $\mu = h_0 + \sigma$. By Proposition 8, we have $\mu \neq \mu^*$. Thus $\mu \notin \mathcal{G}(g)$. \square

A slightly stronger version of Theorem 1 is the following.

Theorem 17. *Given g , let $h_0 \in L^1(\Omega)$ and $K_0 \subset \Omega$ be given by Proposition 8. Let σ be a nonnegative diffuse measure supported in K_0 . If σ is good, then*

$$(8.22) \quad \|\sigma\|_{\mathcal{M}} < \|h_0\|_{L^1}.$$

Proof. Assume σ is good. By Proposition 8, $(h_0 + \sigma)^* = h_0$. Recall that by Theorem 5, $(h_0 + \sigma)^*$ is the closest good measure to $h_0 + \sigma$. Thus,

$$\|\sigma\|_{\mathcal{M}} = \|(h_0 + \sigma) - (h_0 + \sigma)^*\|_{\mathcal{M}} < \|(h_0 + \sigma) - \sigma\|_{\mathcal{M}} = \|h_0\|_{L^1}. \quad \square$$

Corollary 4. *Given g , there exists a diffuse measure $\mu \geq 0$ such that $\varepsilon\mu$ is not good for any $\varepsilon > 0$.*

Proof. Using Remark 3, we can take sequences of disjoint compact sets (K_j) in Ω and L^1 -functions (h_j) , such that each pair K_j, h_j satisfies the assumptions of Proposition 8 and

$$\|h_j\|_{L^1} \leq 1/4^j.$$

Let $h = \sum_j h_j \in L^1(\Omega)$. For each $j \geq 1$, we fix a diffuse measure $\sigma_j \geq 0$ concentrated on K_j such that $\|\sigma_j\|_{\mathcal{M}} = 1/2^j$. Let $\mu = \sum_j \sigma_j \in \mathcal{M}(\Omega)$. Assume, by way of contradiction, that $\varepsilon\mu$ is good for some $\varepsilon > 0$. Since

$$\varepsilon\mu \geq \varepsilon\sigma_j \quad \forall j \geq 1,$$

then $\varepsilon\sigma_j$ is also good. By Theorem 17, this gives

$$\varepsilon < \frac{\|h_j\|_{L^1}}{\|\sigma_j\|_{\mathcal{M}}} \leq \frac{1}{2^j} \quad \forall j \geq 1.$$

Letting $j \rightarrow +\infty$, we get a contradiction. \square

Imposing an additional assumption on the nonlinearity g , one can construct a measure μ of the form $\mu = \theta \mathcal{H}^\alpha \llcorner_K$, for some $\alpha > N - 2$, such that $\mu \notin \mathcal{G}(g)$. To this purpose, one first needs a slight modification of Lemma 10.

Lemma 11. *Assume g is given by*

$$(8.23) \quad g(t) = \frac{1}{(1-t)^{(2-\beta)/\beta}} - 1, \quad t \in [0, 1),$$

where $\beta \in (0, 2)$. Then, for any $\alpha \in (0, \beta)$, there exists $\tilde{v} \in C_0(\overline{\Omega})$ such that

$$(8.24) \quad \Delta \tilde{v} \in L^1(\Omega), \quad \tilde{v} \leq 1 \text{ in } \Omega, \quad \mathcal{H}^{N-2+\alpha}([\tilde{v} = 1]) \in (0, \infty) \quad \text{and} \quad g(\tilde{v}) \in L^1(\Omega).$$

Proof. We adapt the proof of Lemma 10. We consider both cases $N \geq 3$ and $N = 2$ simultaneously. Let \tilde{v} be given by (8.7). Using the same notation as before, let

$$\alpha_j = a_m 2^{-mNj},$$

where

$$(8.25) \quad \frac{\alpha}{N-2+\alpha} < m < \frac{(2-\alpha)\beta}{(2-\beta)\alpha} \frac{\alpha}{N-2+\alpha}$$

and the constant a_m is chosen so that (8.8) holds. Observe that the range of admissible m given by (8.25) is nonempty since $0 < \alpha < \beta < 2$.

Next, let $k_j = j$ and

$$(8.26) \quad \ell_k = 2^{-Nk/(N-2+\alpha)} \quad k \geq 1.$$

With (ℓ_k) defined as above, one can show (see, e.g., [18]) that

$$\mathcal{H}^{N-2+\alpha}(F) \in (0, \infty),$$

where $F = \bigcap_j F_j = [\tilde{v} = 1]$. We now prove (8.11) (or, equivalently, (8.20) if $N = 2$). Note that with our choices of (α_j) and (ℓ_k) , the left-hand side of (8.11) reduces to

$$\sum_{j=1}^{\infty} 2^{mNj \frac{2-\beta}{\beta}} 2^{Nj} 2^{-\frac{N^2 j}{N-2+\alpha}} = \sum_{j=1}^{\infty} 2^{Nj \left(m \frac{2-\beta}{\beta} - \frac{2-\alpha}{N-2+\alpha} \right)},$$

which is finite, by (8.25). First assume $N \geq 3$. Then (8.10) becomes

$$\sum_{j=1}^{\infty} 2^{-mNj} 2^{Nj} 2^{-\frac{N(N-2)j}{N-2+\alpha}} = \sum_{j=1}^{\infty} 2^{-Nj \left(m - \frac{\alpha}{N-2+\alpha} \right)} < \infty,$$

which clearly holds in view of (8.25). Similarly, if $N = 2$, one easily checks that (8.19) is satisfied. Proceeding as in the proof of Lemma 10, we conclude that (8.24) holds. \square

As a consequence, we have the following theorem.

Theorem 18. *Given $\beta \in (0, 2)$, let g be given by (8.23). Then, for any $\alpha \in (0, \beta)$, there exist $\theta_0 > 0$ and $K \subset \Omega$ compact, $\mathcal{H}^{N-2+\alpha}(K) \in (0, \infty)$, such that*

$$(8.27) \quad \theta \mathcal{H}^{N-2+\alpha} \lfloor_K \in \mathcal{G}(g) \quad \text{implies} \quad \theta < \theta_0.$$

Proof. Let

$$\tilde{h}_0 = [-\Delta \tilde{v} + g(\tilde{v})]^+ \quad \text{and} \quad K = [\tilde{v} = 1],$$

where \tilde{v} is given by Lemma 11 above. Proceeding as in the proof of Proposition 8, we have

$$(\tilde{h}_0 + \theta \mathcal{H}^{N-2+\alpha} \lfloor_K)^* = \tilde{h}_0 \quad \text{for all } \theta > 0.$$

Therefore, if $\theta \mathcal{H}^{N-2+\alpha} \lfloor_K$ is good, then as in the proof of Theorem 17, we conclude that

$$\theta \mathcal{H}^{N-2+\alpha}(K) < \|\tilde{h}_0\|_{L^1}.$$

In other words, (8.27) holds with $\theta_0 = \|\tilde{h}_0\|_{L^1} / \mathcal{H}^{N-2+\alpha}(K)$. \square

9 Further properties of μ^* and \mathcal{G}

In this section, we prove some properties of the reduced measures, which should be compared with those in [4]. We start by showing that the reduced measure μ^* need not be the largest good measure $\leq \mu$, contrary to what happens when g is everywhere defined. In fact, we have

Proposition 9. *There exists $\mu \in \mathcal{M}(\Omega)$, $\mu \geq 0$, for which the set*

$$(9.1) \quad \{\lambda \in \mathcal{G} : \lambda \leq \mu\}$$

has no largest element.

Proof. Let K_0, h_0 be given by Proposition 8. Let σ be the capacity measure associated to K_0 . In particular, σ is a nonnegative measure concentrated on K_0 ; moreover, σ is good (see Proposition 7). Let $\mu = h_0 + \sigma$. By Proposition 8, $\mu \notin \mathcal{G}(g)$. Assume, by way of contradiction, that the set given by (9.1) has a largest element, say $\nu \leq \mu$. Clearly, $\nu \geq h_0$ and $\nu \geq \sigma$. Thus,

$$\nu \geq \sup\{h_0, \sigma\} = h_0 + \sigma = \mu.$$

Thus $\nu = \mu$, so μ is a good measure. This is a contradiction. \square

The argument above can be used to establish the following results (in what follows, σ is the capacity measure associated to K_0 , with K_0 and h_0 being given by Proposition 8).

Proposition 10. *There exist good measures $\mu, \nu \geq 0$ such that $\sup\{\mu, \nu\}$ is not good.*

Proof. Take $\mu = h_0, \nu = \sigma$ and use Proposition 8. \square

Proposition 11. *There exist diffuse measures $\mu, \nu \geq 0$ such that $\nu \leq \mu$ but $\mu^* - \nu^*$ is not ≥ 0 .*

Proof. Take $\mu = h_0 + \sigma$ and $\nu = \sigma$. \square

Similarly, the mapping $\mu \mapsto \mu^*$ is not a contraction.

Proposition 12. *There exist diffuse measures $\mu, \nu \geq 0$ such that*

$$\|\mu - \nu\|_{\mathcal{M}} < \|\mu^* - \nu^*\|_{\mathcal{M}}.$$

Proof. Take $\mu = h_0 + \sigma$ and $\nu = \sigma$. \square

We conclude with the following.

Proposition 13. *The set \mathcal{G} is not convex.*

Proof. By Theorem 1, there exists μ diffuse such that $\mu \notin \mathcal{G}$. Applying Theorem 3 in [4], we can decompose μ as

$$\mu = f + \Delta v,$$

where $f \in L^1(\Omega)$, $v \in H_0^1(\Omega) \cap C(\overline{\Omega})$ and $\|v\|_{L^\infty} \leq \frac{1}{3}$. In particular, $2f \in \mathcal{G}$ and $\Delta(2v) \in \mathcal{G}$; however,

$$\frac{2f + \Delta(2v)}{2} = \mu \notin \mathcal{G}. \quad \square$$

Acknowledgments. The second author (A.C.P.) was supported by the NSF grant DMS-0111298 and Sergio Serapioni, Honorary President of Società Trentina Lieviti—Trento (Italy). A.C.P. gratefully acknowledges the invitation to and the warm hospitality of the Math Department at the University of Rome 1. The third author (A.P.) warmly thanks H. Brezis for the invitation to Rutgers University. Part of this work was carried out during these two occasions.

REFERENCES

- [1] A. Ancona, *Une propriété d'invariance des ensembles absorbants par perturbation d'un opérateur elliptique*, Comm. Partial Differential Equations **4** (1979), 321–337.
- [2] L. Boccardo, *On the regularizing effect of strongly increasing lower order terms*. J. Evol. Equ. **3** (2003), 225–236.
- [3] L. Boccardo, T. Gallouët and L. Orsina, *Existence and uniqueness of entropy solutions for nonlinear elliptic equations with measure data*, Ann. Inst. H. Poincaré Anal. Non Linéaire **13** (1996), 539–551.
- [4] H. Brezis, M. Marcus and A. C. Ponce, *Nonlinear elliptic equations with measures revisited*, Annals of Math. Studies, Princeton University Press, to appear. Part of the results were announced in a note by the same authors: *A new concept of reduced measure for nonlinear elliptic equations*, C. R. Acad. Sci. Paris, Ser. I **339** (2004), 169–174.
- [5] H. Brezis and A. C. Ponce, *Remarks on the strong maximum principle*, Differential Integral Equations **16** (2003), 1–12.
- [6] H. Brezis and A. C. Ponce, *Kato's inequality when Δu is a measure*, C. R. Math. Acad. Sci. Paris, Ser. I **338** (2004), 599–604.
- [7] H. Brezis and W. A. Strauss, *Semilinear second-order elliptic equations in L^1* , J. Math. Soc. Japan **25** (1973), 565–590.
- [8] L. Carleson, *Selected Problems on Exceptional Sets*, Van Nostrand, Princeton, 1967.
- [9] G. Dal Maso, F. Murat, L. Orsina and A. Prignet, *Renormalized solutions of elliptic equations with general measure data*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), 741–808.

Author:
Please update

- [10] P. Dall’Aglia and G. Dal Maso, *Some properties of the solutions of obstacle problems with measure data*, Ricerche Mat. **48** (1999), 99–116. Papers in memory of Ennio De Giorgi.
- [11] P. Dall’Aglia and C. Leone, *Obstacles problems with measure data and linear operators*, Potential Anal. **17** (2002), 45–64.
- [12] C. De La Vallée Poussin, *Sur l’intégrale de Lebesgue*, Trans. Amer. Math. Soc. **16** (1915), 435–501.
- [13] C. Dellacherie and P.-A. Meyer, *Probabilités et potentiel*, Chapitres I à IV, Publications de l’Institut de Mathématique de l’Université de Strasbourg, No. XV, Actualités Scientifiques et Industrielles, No. 1372, Hermann, Paris, 1975.
- [14] L. Dupaigne and A. C. Ponce, *Singularities of positive supersolutions in elliptic PDEs*, Selecta Math. (N.S.) **10** (2004), 341–358.
- [15] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.
- [16] M. Fukushima, K. Sato and S. Taniguchi, *On the closable parts of pre-Dirichlet forms and the fine supports of underlying measures*, Osaka J. Math. **28** (1991), 517–535.
- [17] M. Grun-Rehomme, *Caractérisation du sous-différentiel d’intégrandes convexes dans les espaces de Sobolev*, J. Math. Pures Appl. **56** (1977), 149–156.
- [18] A. C. Ponce, *How to construct good measures*, in *Elliptic and Parabolic Equations* (C. Bandle, H. Berestycki, B. Brighi, A. Brillard, M. Chipot, J.-M. Coron, C. Sbordone, I. Shafrir, V. Valente and G. Vergara-Caffarelli, eds.), Progress in Nonlinear Differential Equations and their Applications, 63, Birkhäuser, Basel, 2005, pp. 335–348. A special tribute to the work of Haïm Brezis.
- [19] G. Stampacchia, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier (Grenoble) **15** (1965), 189–258.
- [20] J. L. Vázquez, *On a semilinear equation in \mathbb{R}^2 involving bounded measures*, Proc. Roy. Soc. Edinburgh Sect. A **95** (1983), 181–202.

Louis Dupaigne

LABORATOIRE AMIÉNOIS DE MATHÉMATIQUE FONDAMENTALE ET APPLIQUÉE
 FACULTÉ DE MATHÉMATIQUES ET D’INFORMATIQUE
 UNIVERSITÉ PICARDIE JULES VERNE
 33, RUE SAINT-LEU
 80039 AMIENS CEDEX 1, FRANCE
 email: louis.dupaigne@u-picardie.fr

Augusto C. Ponce

LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE
 FÉDÉRATION DENIS POISSON
 UNIVERSITÉ FRANÇOIS RABELAIS
 37200 TOURS, FRANCE
 email: ponce@lmpt.univ-tours.fr

Alessio Porretta

DIPARTIMENTO DI MATEMATICA
 UNIVERSITÀ DI ROMA “TOR VERGATA”
 VIA DELLA RICERCA SCIENTIFICA 1
 00133 ROMA, ITALY
 email: porretta@mat.uniroma2.it

(Received May 15, 2005)